

High-order short-time expansions for ATM option prices under a tempered stable Lévy model

José E. Figueroa-López*

Ruoting Gong†

Christian Houdré‡

August 29, 2012

Abstract

The short-time asymptotic behavior of option prices for a variety of models with jumps has received much attention in recent years. In the present work, a novel second-order approximation for ATM option prices under an exponential tempered stable model, a rich class of Lévy processes with desirable features for financial modeling, is derived and, then, extended to a model with an additional independent Brownian component. Our method of proof is based on an integral representation of the option price involving the tail probability of the log-return process under the share measure and a suitable change of probability measure under which the process becomes stable. Our approach is sufficiently general to cover a wide class of Lévy processes which satisfy the latter property and whose Lévy densities can be “closely” approximated by a stable density near the origin. The results hereafter shed new light on the connection between both the volatility of the continuous component and the jump parameters and the behavior of ATM option prices near expiration. In the presence of an additional Brownian component, the second-order term, in time- t , is of the form $d_2 t^{(3-Y)/2}$, with the coefficient d_2 depending only on the overall jump intensity of the process and the tail-heaviness parameter Y . This extends the known result that the leading term is $\sigma t^{1/2}/\sqrt{2\pi}$, where σ is the volatility of the continuous component. In contrast, under a pure-jump tempered stable model, the dependence on the overall jump intensity and Y is already reflected in the leading term, which is of the form $d_1 t^{1/Y}$. The information on the relative frequency of negative and positive jumps appears only in the second-order term, which is shown to be of the form $d_2 t$ and whose order of decay turns out to be independent of Y . The asymptotic behavior of the corresponding Black-Scholes implied volatilities is also addressed. Our numerical results show that first-order term typically exhibits rather poor performance and that the second-order term significantly improves the approximation’s accuracy.

AMS 2000 subject classifications: 60G51, 60F99, 91G20, 91G60.

Keywords and phrases: Exponential Lévy models; CGMY and tempered stable models; short-time asymptotics; at-the-money option pricing; implied volatility.

1 Introduction

It is generally recognized that the standard Black-Scholes option pricing model is inconsistent with options data, while remaining a widely used model in practice because of its simplicity. Exponential Lévy models generalize the classical Black-Scholes setup by allowing jumps in stock prices while preserving the independence and stationarity of returns. There are several reasons for introducing jumps in financial modeling. First of all, sudden sharp shifts in the price level of financial assets often occur in practice, and this “jump” risk is quite hard to handled within continuous-paths models. Second, historical asset prices exhibit distributions with so-called stylized features, such as heavy tails, high kurtosis, volatility clustering and leverage effects, which are hard to replicate within purely-continuous frameworks. Finally,

*Department of Statistics, Purdue University, West Lafayette, IN, 47907, USA (figueroa@purdue.edu). Research supported in part by the NSF Grants: DMS-1149692, DMS-0906919.

†Department of Mathematics, Rutgers, The State University of New Jersey, Piscataway, NJ, 08904, USA (rgong@math.rutgers.edu).

‡School of Mathematics, Georgia Institute of Technology, Atlanta, GA, 30332, USA (houdre@math.gatech.edu).

market prices of vanilla options exhibit skewed implied volatilities (relative to changes in the strikes), in contrast to the classical Black-Scholes model which predicts a flat implied volatility smile. Moreover, the fact that the implied volatility smile and skewness phenomenon becomes much more pronounced for short maturities is a clear indication of the presence of jumps. We refer the reader to Cont and Tankov [8] for further motivations on the use of jump processes in financial modeling.

One of the first applications of jump processes in financial modeling is due to Mandelbrot [29], who suggested a pure-jump stable Lévy process Z to model power-like tails and self-similar behavior in cotton price returns. Merton [31] and Press [34] subsequently considered option pricing and hedging problems under an exponential compound Poisson process with Gaussian jumps and an additive independent non-zero Brownian component. A similar exponential compound Poisson jump-diffusion model was more recently studied in Kou [25], where the jump sizes are distributed according to an asymmetric Laplace law. For infinite activity exponential Lévy models, Barndorff-Nielsen [1] introduced the normal inverse Gaussian (NIG) model, while the extension to the generalized hyperbolic class was studied by Eberlein, Keller and Prause [10]. Madan and Seneta [28] introduced the symmetric variance gamma (VG) model while its asymmetric extension was later analyzed in Madan and Milne [27] and Madan, Carr and Chang [26]. Both models are built on Brownian subordination; the main difference being that the log-return process in the NIG model is an infinite variation process with stable like ($\alpha = 1$) behavior of small jumps, while in the VG model the log-price is of finite variation with infinite but relatively low activity of small jumps. The class of “truncated stable” processes was first introduced by Koponen [24]. It was advocated in financial modeling by Cont, Bouchaud and Potters [7] and Matacz [30], and further developed in the context of asset price modeling by Carr, Geman, Madan and Yor [5], who introduced the terminology CGMY. Nowadays, the CGMY model is considered to be a prototype of the general class of models with jumps and enjoys widespread applicability.

The CGMY model is a particular case of the more general KoBoL class of Boyarchenko and Levendorskii [4], which in turn is a subclass of the semi parametric class of tempered stable processes (TSP) introduced by Rosiński [39]. TSP form a rich class with appealing features for financial modeling. One of their most interesting features lies in their short-time and long-time behaviors. Denoting by $Y \in (0, 2)$ the index of a TSP, $L := (L_t)_{t \geq 0}$, it follows that

$$(h^{-1}L_{ht})_{t \geq 0} \xrightarrow{\mathcal{D}} (Z_t)_{t \geq 0}, \quad (h \rightarrow 0), \quad (1.1)$$

where $Z := (Z_t)_{t \geq 0}$ is a strictly α -stable Lévy process with parameter $\alpha = Y$. In fact, (1.1) holds when $Y \in (0, 1)$ provided L is driftless, while it holds when $Y \in (1, 2)$ regardless of the “center” or mean of L . Roughly, (1.1) suggests that the short-time behavior of TSP is akin to that of a stable process, with its heavy-tailed distribution and self-similarity properties which are desirable for financial modeling as emphasized, for example, by Mandelbrot [29]. In contrast,

$$(h^{-1}L_{ht})_{t \geq 0} \xrightarrow{\mathcal{D}} (W_t)_{t \geq 0}, \quad (h \rightarrow \infty),$$

where $(W_t)_{t \geq 0}$ is a Brownian Motion, which suggests that in the long-horizon, the process is Brownian-like. In terms of increments, if we were to consider the consecutive “high-frequency” increments of the process L , say $\{L_{\underline{h}i} - L_{\underline{h}(i-1)}\}_{i \geq 1}$ with $\underline{h} \ll 1$, these will exhibit statistical features consistent with those of a selfsimilar stable time series. But, low-frequency increments, say $\{L_{\bar{h}i} - L_{\bar{h}(i-1)}\}_{i \geq 1}$ with $\bar{h} \gg \underline{h}$, will have Gaussian like distributions.

Stemming in part from its importance for model calibration and testing, small-time asymptotics of option prices have received a lot of attention in recent years (see, e.g., [2], [3], [11], [12], [13], [17], [18], [19], [20], [21], [33], [35], [43]). We shall review here only the studies most closely related to ours, focusing in particular on the at-the-money (ATM) case. Carr and Wu [9] first analyzed, partially via heuristic arguments, the first order asymptotic behavior of an Itô semimartingale with jumps. Concretely, ATM option prices of pure-jump models of bounded variation decrease at the rate $O(t)$, while they are just $O(t^{1/2})$ under the presence of a Brownian component. By considering a stable pure-jump component, [9] also showed that, in general, the rate could be $O(t^{\beta})$, for some $\beta \in (0, 1)$. Muhle-Karbe and Nutz [32] formally showed that, under the presence of a continuous-time component, the leading term of ATM option prices is of order $\sqrt{t}$, for a relatively general class of Itô models, while for a more general type of Itô processes with α -stable-like small jumps, the leading term is $O(t^{1/\alpha})$ (see also [13, Proposition 4.2], [15, Theorem 3.7], and [43, Proposition 5] for related results in exponential Lévy models). However, none of these papers have obtained second or higher order asymptotics for the ATM option prices, which are arguably more relevant for calibration purposes, given that the most liquid options are of this type.

In the present paper, we study the small-time behavior for at-the-money call (or equivalently, put) option prices

$$\mathbb{E}(S_t - S_0)^+ = S_0 \mathbb{E}(e^{X_t} - 1)^+, \quad (1.2)$$

under the exponential Lévy model

$$S_t := S_0 e^{X_t}, \quad (1.3)$$

where $X := (X_t)_{t \geq 0}$ is the superposition of a *proper tempered stable* Lévy process $L := (L_t)_{t \geq 0}$ in the sense of [39] (see Section 2 below for a precise definition) and of an independent Brownian motion $(\sigma W_t)_{t \geq 0}$; i.e.,

$$X_t := \sigma W_t + L_t, \quad (1.4)$$

where $W := (W_t)_{t \geq 0}$ is a standard Brownian motion independent of L . Throughout, x^+ is the positive part of x . As it will be shown below, the first order asymptotic behavior of (1.2) in short-time under (1.4) takes the form:

$$\lim_{t \rightarrow 0} t^{-1/Y} \mathbb{E}(S_t - S_0)^+ = S_0 \mathbb{E}(Z^+), \quad (1.5)$$

where Z is a centered α -stable random variable with $\alpha = Y$ under $\mathbb{P}$. When $\sigma \neq 0$, $Z \sim \mathcal{N}(0, \sigma^2)$ (i.e., $\alpha = 2$) and, thus, $\mathbb{E}(Z^+) = \sigma/\sqrt{2\pi}$, as already shown in [43] and [35]. When $\sigma = 0$ and L is symmetric, the characteristic function of Z is explicitly given by

$$\mathbb{E}e^{iuZ} = e^{-2C\Gamma(-Y)|\cos(\frac{1}{2}Y\pi)||u|^Y},$$

where C is a measure of the overall jump-intensity of the process. In that case, (see (25.6) in [42]),

$$d_1 := \mathbb{E}(Z^+) = \frac{1}{\pi} \Gamma\left(1 - \frac{1}{Y}\right) \left(2C\Gamma(-Y) \left|\cos\left(\frac{\pi Y}{2}\right)\right|\right)^{1/Y}. \quad (1.6)$$

We refer the reader to Remark 3.5 below for the explicit formula for the leading term in the general case. Interestingly enough, under the presence of a continuous component ($\sigma \neq 0$), the first-order asymptotic term only reflects information on the continuous-time volatility, in sharp contrast with the pure-jump case where the leading term depends on the overall jump-intensity parameter C and the index Y , which in turn controls the tail-heaviness of the distributions.

The asymptotic formula (1.5) is in agreement with the result of Tankov [43] (see Theorem 5 therein), who showed that for a pure-jump Lévy process, whose characteristic function is similar to that of a stable process, the rate of convergence of ATM option prices is of order $t^{1/Y}$. Concretely, it is shown in [43] that if the characteristic function of the Lévy process $(X_t)_{t \geq 0}$ is such that

$$\mathbb{E}(e^{iuX_1}) = \exp(iu\gamma - |u|^Y f(u)), \quad (1.7)$$

for $1 < Y < 2$ and a function f satisfying $\lim_{u \nearrow \infty} f(u) = \hat{c}_+$ and $\lim_{u \searrow -\infty} f(u) = \hat{c}_-$ for some constants $0 < \hat{c}_+, \hat{c}_- < \infty$, then

$$\lim_{t \rightarrow 0} t^{-1/Y} \mathbb{E}(S_t - S_0)^+ = S_0 \frac{1}{2\pi} \left(\hat{c}_+^{1/Y} + \hat{c}_-^{1/Y}\right). \quad (1.8)$$

As it turns out, the characteristic function of a tempered stable process satisfies the specification (1.7) (see Theorem 2.9 in [39]) and, thus, (1.8) can be applied to explicitly write the leading order term in (1.5). Alternatively, we can also use a result of [32] (see Theorem 4.4 therein), where a formula similar to (1.5) is shown for a more general class of pure-jump martingales. Nevertheless, as previously mentioned, none of these papers obtained second or higher order terms for the ATM option prices and the approaches therein do not seem amenable to study higher-order asymptotics.

The main result of the present paper establishes a second order correction term for the approximation (1.5) in a general tempered stable Lévy model with a possible nonzero independent Brownian component. To the best of our knowledge, these are the first results of this kind in the literature of exponential Lévy models. We show that the second-order asymptotic behavior of the ATM call option price (1.2) in short-time is of the form

$$\frac{1}{S_0} \mathbb{E}(S_t - S_0)^+ = d_1 t^{\frac{1}{Y}} + d_2 t + o(t), \quad (t \rightarrow 0),$$

in the pure-jump tempered stable Lévy case (i.e., $\sigma = 0$), while in the presence of a non-zero independent Brownian component (i.e., $\sigma \neq 0$),

$$\frac{1}{S_0} \mathbb{E}(S_t - S_0)^+ = d_1 t^{\frac{1}{2}} + d_2 t^{\frac{3-Y}{2}} + o\left(t^{\frac{3-Y}{2}}\right), \quad (t \rightarrow 0),$$

for different constants d_1 and d_2 that we will determine explicitly. To wit, we found that, under the presence of a nonzero Gaussian component, the second-order term depends only on the overall jump intensity parameter C and the tail-heaviness parameter Y . However, for a pure-jump case, the tail features of the Lévy measure are already present in the second-order term. The above asymptotic behaviors should also be compared to the corresponding behavior under the standard Black-Scholes model, where it is known that (see, e.g., [19, Corollary 3.4])

$$\mathbb{E}(e^{\sigma W_t - \frac{\sigma^2}{2}t} - 1)^+ = \frac{\sigma}{\sqrt{2\pi}} t^{\frac{1}{2}} - \frac{\sigma^3}{24\sqrt{2\pi}} t^{\frac{3}{2}} + O(t^{\frac{5}{2}}).$$

As a byproduct of our asymptotic results for option prices, we also give the asymptotic behavior of the corresponding Black-Scholes implied volatilities.

The derivation of our second-order approximation results builds on two facts. First, as in [13], we make use of the following representation of Carr and Madan [6]:

$$\frac{1}{S_0} \mathbb{E}(S_t - S_0)^+ = \mathbb{P}^*(X_t > E) = \int_0^\infty e^{-x} \mathbb{P}^*(X_t > x) dx, \quad (1.9)$$

where $\mathbb{P}^*$ is the martingale probability measure obtained when one takes the stock as the numéraire (i.e., $\mathbb{P}^*(A) := \mathbb{E}(S_t \mathbf{1}_A)$) and E is an independent mean-one exponential random variable under $\mathbb{P}^*$. The measure $\mathbb{P}^*$ is also called the share measure (see [6]). As it turns out, under $\mathbb{P}^*$, $(X_t)_{t \geq 0}$ admits a decomposition similar to (1.4),

$$X_t = \sigma W_t^* + L_t^*, \quad t \geq 0, \quad (1.10)$$

where $W^* := (W_t^*)_{t \geq 0}$ is a Wiener process while $L^* := (L_t^*)_{t \geq 0}$ is a proper tempered stable process, independent of W^* . Second, we change probability measures from $\mathbb{P}^*$ to another probability measure, say $\tilde{\mathbb{P}}$, under which $(L_t^*)_{t \geq 0}$ is a stable Lévy process and $(W_t^*)_{t \geq 0}$ is still a standard Brownian motion independent of L^* . This change will in turn enable us to exploit some key features of stable processes such as self-similarity and the tail behavior of their marginal densities.

As explained above, the class of tempered stable Lévy processes is quite rich and enjoys several desirable properties for financial modeling. A natural question is whether our approach is general enough to handle any Lévy process, which satisfies the fundamental property of being stable under a suitable change of probability measure. As a byproduct of our analysis (see Remark 4.4 for precise arguments), we found this to be the case under a key additional condition implying that the Lévy density of the process can be “closely” approximated by a stable density near the origin. Concretely, denoting the Lévy density of the process under the risk-neutral measure by s , our results will hold true (modulo some additional mild conditions) if, for some $Y \in (1, 2)$, the function $q(x) := s(x)/|x|^{Y+1}$ is such that

$$\lim_{x \searrow 0} \frac{1}{x} (C_+ - q(x)) = M, \quad \lim_{x \nearrow 0} \frac{1}{x} (C_- - q(x)) = -G, \quad (1.11)$$

for some constants $C_+, C_-, G > 0$ and $M > 1$.

The present article is organized as follows. Section 2 contains preliminary definitions and properties related to tempered stable processes and probability measure transformations, which will be needed throughout the paper. Section 3 establishes the second-order asymptotics of the call option price under the pure-jump tempered stable model ($\sigma = 0$). Section 4 establishes the second-order asymptotics of the call option price under the tempered stable model with an additional independent non-zero Brownian component ($\sigma \neq 0$). Section 5 illustrates our second-order asymptotics for the important particular class of CGMY models, recovering our preliminary results first presented in [16]. Section 6 assesses the performance of the asymptotic expansions through a detailed numerical analysis for the CGMY model. The proofs of our main results are deferred to the Appendices.

2 Tempered stable processes

Let $L := (L_t)_{t \geq 0}$ be a pure-jump Lévy process with triplet $(0, b, \nu)$ and let $W := (W_t)_{t \geq 0}$ be a Wiener process, independent of L , defined on a complete filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathbb{P})$. We assume zero interest rate and that $\mathbb{P}$ is a martingale measure for the exponential Lévy model

$$S_t = S_0 e^{X_t}, \quad \text{with} \quad X_t := \sigma W_t + L_t,$$

where $(S_t)_{t \geq 0}$ represents the price process of a non-dividend paying risky asset. Equivalently, the Lévy triplet (σ^2, b, ν) of $X := (X_t)_{t \geq 0}$ is such that

$$1 = \mathbb{E}(e^{X_t}) = \exp \left(t \left[bu + \frac{\sigma^2}{2} + \int_{\mathbb{R}_0} (e^x - 1 - x \mathbf{1}_{\{|x| \leq 1\}}) \nu(dx) \right] \right), \quad (t \geq 0), \quad (2.1)$$

where hereafter $\mathbb{R}_0 := \mathbb{R} \setminus \{0\}$ and the Lévy triplets are given relative to the truncation function $\mathbf{1}_{\{|x| \leq 1\}}$ (see Section 8 in [42]). Without loss of generality, we also assume that $(X_t)_{t \geq 0}$ is the canonical process $X_t(\omega) := \omega(t)$ defined on the canonical space $\Omega = \mathbb{D}([0, \infty), \mathbb{R})$ (the space of càdlàg functions $\omega : [0, \infty) \rightarrow \mathbb{R}$) equipped with the σ -field $\mathcal{F} := \sigma(X_s : s \geq 0)$ and the right-continuous filtration $\mathcal{F}_t := \cap_{s > t} \sigma(X_u : u \leq s)$.

In this paper, we consider an exponential tempered stable Lévy model, i.e., throughout we assume that, under $\mathbb{P}$, $L := (L_t)_{t \geq 0}$ is a *proper tempered stable* Lévy process in the sense of [39] with index $Y \in (1, 2)$. That is, the Lévy measure ν of L under $\mathbb{P}$ admits a density $s : \mathbb{R}_0 \rightarrow [0, \infty)$ of the form (see (2.7) in [39]):

$$s(x) = |x|^{-Y-1} q(x),$$

with a function $q : \mathbb{R} \setminus \{0\} \rightarrow [0, \infty)$ given by

$$q(x) := \begin{cases} q^+(x) & x > 0, \\ q^-(-x) & x < 0, \end{cases} := \begin{cases} \int_0^\infty e^{-x/s} s^Y \rho(ds) & x > 0, \\ \int_{-\infty}^0 e^{-x/s} |s|^Y \rho(ds) & x < 0, \end{cases} \quad (2.2)$$

for a nonzero measure ρ on $\mathbb{R}_0$ such that

$$\int_{\mathbb{R}_0} |s|^Y \rho(ds) < \infty.$$

By Bernstein's theorem, the representation (2.2) is equivalent to asking that $q^+ : \mathbb{R}_+ \rightarrow [0, \infty)$ and $q^- : \mathbb{R}_+ \rightarrow [0, \infty)$ are completely monotone functions, meaning that $(-1)^n \frac{\partial^n}{\partial x^n} q^\pm(x) > 0$, for all $x > 0$. In particular, $q^\pm : \mathbb{R}_+ \rightarrow [0, \infty)$ are strictly decreasing and convex. A prototypical tempered stable process is the CGMY process (also called the KoBoL model) of widespread applicability in mathematical finance. For the CGMY process, $\rho(ds) := CM^Y \delta_{\{M^{-1}\}}(ds) + CG^Y \delta_{\{-G^{-1}\}}(ds)$ so that the Lévy measure of L is given by

$$\nu(dx) = \left(\frac{C e^{-Mx}}{x^{1+Y}} \mathbf{1}_{\{x > 0\}} + \frac{C e^{Gx}}{|x|^{1+Y}} \mathbf{1}_{\{x < 0\}} \right) dx. \quad (2.3)$$

For $\mathbb{P}$ to be a martingale measure, we need that $\mathbb{E}e^{X_t} < \infty$ or, equivalently, that

$$\int_1^\infty e^x \nu(dx) = \int_1^\infty e^x q(x) x^{-Y-1} dx < \infty. \quad (2.4)$$

It is not hard to see that a necessary and sufficient condition for (2.4) to hold is that

$$\rho((1, \infty)) = 0. \quad (2.5)$$

Indeed, suppose that (2.4) holds. Then, for any $s_0 > 1$, we will have that

$$\int_1^\infty e^x q(x) x^{-Y-1} dx \geq \int_1^\infty x^{-Y-1} \int_{s_0}^\infty e^{-x(\frac{1}{s}-1)} s^Y \rho(ds) dx \geq \int_1^\infty x^{-Y-1} e^{-x(\frac{1}{s_0}-1)} dx \int_{s_0}^\infty s^Y \rho(ds) = \infty,$$

unless $\rho([s_0, \infty)) = 0$, for any $s_0 > 1$. On the other hand, if (2.5) is satisfied, $q(x) = \int_0^1 e^{-x/s} s^Y \rho(ds) \leq e^{-x} \int_0^1 s^Y \rho(ds)$ and, thus,

$$\int_1^\infty e^x \nu(dx) = \int_1^\infty e^x q(x) x^{-Y-1} dx \leq \int_0^1 s^Y \rho(ds) \int_1^\infty x^{-Y-1} dx < \infty.$$

Therefore, from now on, we shall enforce the condition (2.5). We also make the following standing assumptions:

Assumption 2.1. The function $q : \mathbb{R}_0 \rightarrow [0, \infty)$ is such that

$$(i) \limsup_{x \searrow 0} -q'(x) < \infty, \quad (ii) \limsup_{x \nearrow 0} q'(x) < \infty, \quad (iii) \limsup_{|x| \rightarrow \infty} \frac{|\ln q(x)|}{|x|} < \infty. \quad (2.6)$$

Since

$$-q'(x) = \int_0^\infty e^{-x/s} s^{Y-1} \rho(ds) \geq 0, \quad \text{for } x > 0, \quad q'(x) = \int_{-\infty}^0 e^{-x/s} |s|^{Y-1} \rho(ds) \geq 0, \quad \text{for } x < 0,$$

conditions (2.6-i) and (2.6-ii) are equivalent to asking that the measure ρ satisfies

$$\int_{\mathbb{R}_0} |s|^{Y-1} \rho(ds) < \infty. \quad (2.7)$$

Hereafter, we set

$$C_+ := q(0^+) = \int_0^\infty s^Y \rho(ds), \quad C_- := q(0^-) = \int_{-\infty}^0 |s|^Y \rho(ds). \quad (2.8)$$

It is also convenient to introduced the following standardized versions of q and ρ :

$$\begin{aligned} \bar{\rho}(dx) &:= \left(\frac{1}{C_+} \mathbf{1}_{\{x>0\}} + \frac{1}{C_-} \mathbf{1}_{\{x<0\}} \right) \rho(dx), \\ \bar{q}(x) &:= \frac{q(x)}{C_+} \mathbf{1}_{\{x>0\}} + \frac{q(x)}{C_-} \mathbf{1}_{\{x<0\}} = \begin{cases} \int_0^\infty e^{-x/s} s^Y \bar{\rho}(ds) & x > 0, \\ \int_{-\infty}^0 e^{-x/s} |s|^Y \bar{\rho}(ds) & x < 0. \end{cases} \end{aligned}$$

Note that these standardizations ensure that

$$\bar{q}(0^+) = \int_0^\infty s^Y \bar{\rho}(ds) = 1, \quad \bar{q}(0^-) = \int_{-\infty}^0 |s|^Y \bar{\rho}(ds) = 1. \quad (2.9)$$

Under (2.9), hereafter $\bar{q}(0) := 1$. The following properties of q are crucial to obtain the second-order asymptotics:

Lemma 2.2. *Let*

$$M := -\bar{q}'(0^+) = \int_0^\infty s^{Y-1} \bar{\rho}(ds), \quad \text{and} \quad G := \bar{q}'(0^-) = \int_{-\infty}^0 |s|^{Y-1} \bar{\rho}(ds).$$

Then, under the conditions (2.6-i) and (2.6-ii),

$$\lim_{\delta \searrow 0} \frac{1}{\delta} (1 - \bar{q}(\delta y)) = yM \mathbf{1}_{\{y>0\}} + |y|G \mathbf{1}_{\{y<0\}}. \quad (2.10)$$

In particular,

$$(i) \lim_{\delta \searrow 0} \bar{q}(\delta y)^{1/\delta} = e^{-yM} \mathbf{1}_{\{y>0\}} + e^{-|y|G} \mathbf{1}_{\{y<0\}}, \quad (ii) \lim_{\delta \searrow 0} \frac{1}{\delta} \ln \bar{q}(\delta y) = -yM \mathbf{1}_{\{y>0\}} - |y|G \mathbf{1}_{\{y<0\}}. \quad (2.11)$$

Proof. Let $y > 0$. Then, for $\delta > 0$,

$$\frac{1}{\delta} (1 - \bar{q}(\delta y)) = \int_0^\infty \frac{1}{\delta} \left(1 - e^{-\frac{\delta y}{s}}\right) s^Y \bar{\rho}(ds) \xrightarrow{\delta \searrow 0} yM,$$

where the limit follows from dominated convergence and the inequality $0 \leq 1 - e^{-\delta y/s} \leq \delta y/s$. Similarly, for $y < 0$ and $\delta > 0$,

$$\frac{1}{\delta} (1 - \bar{q}(\delta y)) = \int_{-\infty}^0 \frac{1}{\delta} \left(1 - e^{-\frac{\delta y}{s}}\right) |s|^Y \bar{\rho}(ds) \xrightarrow{\delta \searrow 0} |y|G.$$

To show (2.11-i), note that

$$\lim_{\delta \searrow 0} \bar{q}(\delta y)^{1/\delta} = \lim_{\delta \rightarrow 0} \left(1 - \delta \left(\frac{1 - \bar{q}(y\delta)}{\delta}\right)\right)^{1/\delta} = e^{-yM} \mathbf{1}_{\{y>0\}} + e^{-|y|G} \mathbf{1}_{\{y<0\}}.$$

Then, (2.11-ii) follows from (2.11-i) by taking logarithms on both sides. $\square$

Following a density transformation construction as given in Sato [42] (see Definition 33.4 and Example 33.14 there) and using the martingale condition $\mathbb{E}e^{X_t} = 1$, we define a probability measure $\mathbb{P}^*$ on $(\Omega, \mathcal{F})$ via

$$\frac{d\mathbb{P}^*|_{\mathcal{F}_t}}{d\mathbb{P}|_{\mathcal{F}_t}} = e^{X_t}, \quad (t \geq 0); \quad (2.12)$$

i.e., $\mathbb{P}^*(B) = \mathbb{E}(e^{X_t} \mathbf{1}_B)$, for any $B \in \mathcal{F}_t$ and $t \geq 0$. The measure $\mathbb{P}^*$ can be interpreted as the martingale measure when using the stock price as the numéraire. Under this probability measure, $(X_t)_{t \geq 0}$ has the representation $(\sigma W_t^* + L_t^*)_{t \geq 0}$ where, under $\mathbb{P}^*$, $W^* := (W_t^*)_{t \geq 0}$ is a Wiener process and $L^* := (L_t^*)_{t \geq 0}$ is a Lévy process with triplet $(0, b^*, \nu^*)$ given by

$$\nu^*(dx) := e^x \nu(dx) = e^x s(x) dx, \quad b^* := b + \int_{|x| \leq 1} x(e^x - 1) s(x) dx + \sigma^2,$$

which is moreover independent of L^* . Hereafter, we set

$$q^*(x) = q(x)e^x, \quad s^*(x) = |x|^{-Y-1} q^*(x), \quad \bar{q}^*(x) := \bar{q}(x)e^x.$$

Remark 2.3. As it turns out, under (2.4) (equivalently, under (2.5)), ν^* corresponds to a tempered stable Lévy measure. Indeed, setting $T(s) := (\frac{1}{s} - 1)^{-1}$ and, for $x > 0$,

$$q^*(x) := e^x q(x) = e^x \int_0^\infty e^{-\frac{x}{s}} s^Y \rho(ds) = \int_0^1 e^{-x(\frac{1}{s}-1)} s^Y \rho(ds) = \int_0^\infty e^{-\frac{x}{u}} u^Y \rho^*(du), \quad (2.13)$$

with $\rho^*(du)|_{(0,\infty)} := \mathbf{1}_{(0,\infty)}(u)(1+u)^{-Y} \rho \circ T^{-1}(du)$, while, for $x < 0$,

$$q^*(x) := e^x q(x) = e^x \int_{-\infty}^0 e^{-x/s} |s|^Y \rho(ds) = \int_{-\infty}^0 e^{-x(\frac{1}{s}-1)} (-s)^Y \rho(ds) = \int_{-\infty}^0 e^{-\frac{x}{u}} (-u)^Y \rho^*(du), \quad (2.14)$$

with $\rho^*(du)|_{(-\infty,0)} := \mathbf{1}_{(-1,0)}(u)(1+u)^{-Y} \rho \circ T^{-1}(du)$.

The following simple consequence of Lemma 2.2 will also be needed in the sequel:

Lemma 2.4. *Let*

$$M^* := M - 1 = -\bar{q}'(0^+) - 1 \quad \text{and} \quad G^* := G + 1 := \bar{q}'(0^-) + 1,$$

and let the conditions (2.4), (2.6-i), and (2.6-ii) be satisfied. Then, $M^* > 0$ and, for any $y \in \mathbb{R}_0$,

$$(i) \lim_{\delta \searrow 0} \bar{q}^*(\delta y)^{1/\delta} = e^{-yM^*} \mathbf{1}_{\{y>0\}} + e^{-|y|G^*} \mathbf{1}_{\{y<0\}}, \quad (ii) \lim_{\delta \searrow 0} \frac{1}{\delta} \ln \bar{q}^*(\delta y) = -yM^* \mathbf{1}_{\{y>0\}} - |y|G^* \mathbf{1}_{\{y<0\}}. \quad (2.15)$$

Proof. The first assertion is clear since, as it was shown above, (2.4) implies (2.5) and, therefore, using (2.9),

$$M^* = M - 1 = \int_0^\infty s^Y \left(\frac{1}{s} - 1 \right) \bar{\rho}(ds) = \int_0^1 s^Y \left(\frac{1}{s} - 1 \right) \bar{\rho}(ds) > 0.$$

Next, from Lemma 2.2,

$$\frac{1}{\delta} (1 - \bar{q}^*(\delta y)) = \frac{1}{\delta} (1 - \bar{q}(\delta y) e^{\delta y}) = \frac{1}{\delta} (1 - \bar{q}(\delta y)) + \frac{\bar{q}(\delta y)}{\delta} (1 - e^{\delta y}) \xrightarrow{\delta \searrow 0} y(M - 1) \mathbf{1}_{y>0} + |y|(G + 1) \mathbf{1}_{y<0}.$$

Using the above and proceeding as in the proof of Lemma 2.2 lead to (2.15). $\square$

An important tool thereafter is to change probability measures from $\mathbb{P}^*$ to another probability measure, say $\tilde{\mathbb{P}}$, under which $(L_t^*)_{t \geq 0}$ is a stable Lévy process and $(W_t^*)_{t \geq 0}$ is a Wiener process independent of L^* . Concretely, let

$$\tilde{\nu}(dx) := |x|^{-Y-1} (C_+ \mathbf{1}_{\{x>0\}} + C_- \mathbf{1}_{\{x<0\}}) dx, \quad \tilde{b} = b^* + \int_{|x| \leq 1} x(\tilde{\nu} - \nu^*)(dx).$$

Note that $\tilde{\nu}$ is the Lévy measure of a Y -stable Lévy process and that

$$\tilde{\nu}(dx) = e^{\varphi(x)} \nu^*(dx), \quad \text{with } \varphi(x) := -\ln \bar{q}^*(x) = -\ln \bar{q}(x) - x.$$

Hence, by virtue of Theorem 33.1 in [42], there exists a probability measure $\tilde{\mathbb{P}}$ locally equivalent¹ to $\mathbb{P}^*$ such that $(X_t)_{t \geq 0}$ is a Lévy process with Lévy triplet $(\sigma^2, \tilde{b}, \tilde{\nu})$ under $\tilde{\mathbb{P}}$. Throughout, $\tilde{\mathbb{E}}$ denotes expectation under $\tilde{\mathbb{P}}$.

Remark 2.5. By virtue of [42, Theorem 33.1], the previous change of probability measure is valid since, under our assumptions, the following condition is satisfied:

$$\int_{\mathbb{R}_0} \left(e^{\varphi(x)/2} - 1 \right)^2 \nu^*(dx) = \int_{\mathbb{R}_0} \left(e^{\varphi(x)/2} - 1 \right)^2 e^{-\varphi(x)} \tilde{\nu}(dx) < \infty.$$

Indeed, the integral above can be expressed as

$$C_+ \int_0^\infty \left(e^{\frac{1}{2} \ln \bar{q}^*(x)} - 1 \right)^2 |x|^{-Y-1} dx + C_- \int_{-\infty}^0 \left(e^{\frac{1}{2} \ln \bar{q}^*(x)} - 1 \right)^2 |x|^{-Y-1} dx, \quad (2.16)$$

and, in light of (2.15), the integrand in the first integral is such that

$$\left(e^{\frac{1}{2} \ln \bar{q}^*(x)} - 1 \right)^2 \sim \frac{1}{4} (\ln \bar{q}^*(x))^2 \sim \frac{(M^*)^2}{4} x^2, \quad \text{as } x \searrow 0.$$

This shows that the first integral is finite on any interval $(0, \varepsilon)$. Outside any neighborhood of the origin, this integral is finite in view of (2.4). The second integral in (2.16) can be handled using again (2.15) and the fact that $\ln \bar{q}^*(x) < 0$ when $x < 0$.

Denoting $\tilde{\gamma} := \tilde{\mathbb{E}}X_1 = \tilde{\mathbb{E}}L_1^*$, we recall that the centered process $(Z_t)_{t \geq 0}$, defined by

$$Z_t := L_t^* - t\tilde{\gamma}, \quad (2.17)$$

is a strictly Y -stable under $\tilde{\mathbb{P}}$ and, thus, is also self-similar; i.e.,

$$(t^{-1/Y} Z_{ut})_{u \geq 0} \stackrel{\mathcal{D}}{=} (Z_u)_{u \geq 0}, \quad (2.18)$$

for any $t > 0$. Let us now recall some standard results on stable Lévy processes. Throughout, let p_Z denote the marginal density function of Z_1 under $\tilde{\mathbb{P}}$. It is well known (see, e.g., (14.37) in [42] and references therein) that

$$p_Z(v) \sim C_\pm |v|^{-Y-1}, \quad \text{as } v \rightarrow \pm\infty, \quad \text{respectively.} \quad (2.19)$$

¹Equivalently, there exists a process $(U_t)_{t \geq 0}$ such that $\tilde{\mathbb{P}}^*(B) = \mathbb{E}^*(e^{U_t} \mathbf{1}_B)$, for $t \geq 0$ and $B \in \mathcal{F}_t$.

In particular, for any $v > 0$, as $t \rightarrow 0$,

$$\tilde{\mathbb{P}}(Z_t \geq v) = \tilde{\mathbb{P}}\left(Z_1 \geq t^{-1/Y}v\right) \sim t\tilde{\nu}([v, \infty)) = t\frac{C_+}{Y}v^{-Y}, \quad (2.20)$$

$$\tilde{\mathbb{P}}(Z_t \leq -v) = \tilde{\mathbb{P}}\left(Z_1 \leq -t^{-1/Y}v\right) \sim t\tilde{\nu}((-\infty, -v]) = t\frac{C_-}{Y}|v|^{-Y}. \quad (2.21)$$

The following tail estimate can also be deduced from (2.19) (see Appendix C for its proof):

$$\tilde{\mathbb{P}}\left(|Z_1| \geq t^{-1/Y}v\right) = \tilde{\mathbb{P}}(|Z_t| \geq v) \leq \kappa t|v|^{-Y}, \quad (2.22)$$

for all $v > 0$ and $0 < t \leq 1$ and some absolute constant $\kappa < \infty$.

Of use in what follows is the following representation of the density process (see Theorem 33.2 in [42])

$$\frac{d\tilde{\mathbb{P}}|_{\mathcal{F}_t}}{d\mathbb{P}^*|_{\mathcal{F}_t}} = e^{U_t}, \quad (2.23)$$

with

$$U_t := \lim_{\epsilon \rightarrow 0} \left(\sum_{\substack{s \leq t: \\ |\Delta X_s| > \epsilon}} \varphi(\Delta X_s) - t \int_{|x| > \epsilon} \left(e^{\varphi(x)} - 1 \right) \nu^*(dx) \right).$$

The process $(U_t)_{t \geq 0}$ can be expressed in terms of the jump-measure $N(dt, dx) := \#\{(s, \Delta X_s) \in dt \times dx\}$ of the process $(X_t)_{t \geq 0}$ and its compensated measure $\bar{N}(dt, dx) := N(dt, dx) - \tilde{\nu}(dx)dt$ (under $\tilde{\mathbb{P}}$). Indeed,

$$\begin{aligned} U_t &= \lim_{\epsilon \rightarrow 0} \left(\int_0^t \int_{|x| > \epsilon} \varphi(x) N(ds, dx) - t \int_{|x| > \epsilon} \left(e^{\varphi(x)} - 1 \right) e^{-\varphi(x)} \tilde{\nu}(dx) \right) \\ &= \lim_{\epsilon \rightarrow 0} \left(\int_0^t \int_{|x| > \epsilon} \varphi(x) \bar{N}(ds, dx) + t \int_{|x| > \epsilon} \left(e^{-\varphi(x)} - 1 + \varphi(x) \right) \tilde{\nu}(dx) \right). \end{aligned}$$

Finally, from the definition of Poisson integrals under compensated Poisson measures as in [23] (see Theorem 10.15 therein), we get

$$U_t = \tilde{U}_t + \eta t, \quad (2.24)$$

with

$$\tilde{U}_t := \int_0^t \int_{\mathbb{R}_0} \varphi(x) \bar{N}(ds, dx), \quad \eta := \int_{\mathbb{R}_0} \left(e^{-\varphi(x)} - 1 + \varphi(x) \right) \tilde{\nu}(dx). \quad (2.25)$$

Remark 2.6. Both $\tilde{U}_t$ and η above are indeed well defined since from (2.4), (2.6-iii), (2.15-ii), and since $\varphi(x) \geq 0$ for all $x \in \mathbb{R}_-$ and $Y \in (1, 2)$,

$$\begin{aligned} \int_{|x| \leq 1} \left| e^{-\varphi(x)} - 1 + \varphi(x) \right| |x|^{-Y-1} dx &\leq C_1 \int_{|x| \leq 1} \varphi^2(x) |x|^{-Y-1} dx \leq C_2 \int_{|x| \leq 1} |x|^{1-Y} dx < \infty, \\ \int_{x < -1} \left| e^{-\varphi(x)} - 1 + \varphi(x) \right| |x|^{-Y-1} dx &\leq C_1 \int_{x < -1} |\varphi(x)| |x|^{-Y-1} dx \leq C_2 \int_{x < -1} |x|^{-Y} dx < \infty, \\ \int_{x > 1} \left| e^{-\varphi(x)} - 1 + \varphi(x) \right| |x|^{-Y-1} dx &\leq C_1 \int_{x > 1} e^{\ln q(x) + x} x^{-Y-1} dx + \int_{x > 1} |-1 + \varphi(x)| x^{-Y-1} dx < \infty, \end{aligned}$$

for some constants $C_1, C_2 < \infty$.

Finally, let us also note the decomposition of the process X in terms of the compensated measure $\bar{N}(dt, dx) := N(dt, dx) - \tilde{\nu}(dx)dt$:

$$X_t = \sigma W_t^* + Z_t + t\tilde{\gamma} = \sigma W_t^* + \int_0^t \int_{\mathbb{R}_0} x \bar{N}(ds, dx) + t\tilde{\gamma}. \quad (2.26)$$

The representation (2.26) can be deduced from the Lévy-Itô decomposition of the process $(X_t)_{t \geq 0}$ (cf. Theorem 13.4 and Corollary 13.7 in [23] or Theorem 19.2 in [42]), the stated Lévy triplet $(0, \tilde{b}, \tilde{\nu})$ of X under $\tilde{\mathbb{P}}$, and since $\tilde{b}$ must be such that $\tilde{\mathbb{E}}(X_t) = \tilde{\gamma}t$, by construction.

3 The pure-jump tempered stable model

In this section, we find the second-order asymptotic behavior for the at-the-money call option prices (1.2) in the pure-jump tempered stable process (i.e. $\sigma = 0$ and $X_t = L_t^*$). The proofs of all the results of this section are deferred to Appendix A. Before stating our first result, we need to rewrite the call option price (1.2) in a suitable form.

Lemma 3.1. *With the probability measure $\tilde{\mathbb{P}}$ defined in (2.23) and the parameter $\tilde{\gamma} := \tilde{\mathbb{E}}X_1 = \tilde{\mathbb{E}}L_1^*$,*

$$t^{-\frac{1}{Y}} \frac{1}{S_0} \mathbb{E}(S_t - S_0)^+ = e^{-(\tilde{\gamma} + \eta)t} \int_{-\tilde{\gamma}t^{1-1/Y}}^{\infty} e^{-t^{1/Y}v} \tilde{\mathbb{E}}\left(e^{-\tilde{U}_t} \mathbf{1}_{\{t^{-1/Y}Z_t \geq v\}}\right) dv. \quad (3.1)$$

The next two lemmas are crucial to obtain the main result of the section:

Lemma 3.2. *In the setting and under the assumptions of Section 2, for any $\xi \geq 0$,*

$$(i) \lim_{t \rightarrow 0} \tilde{\mathbb{E}}\left(e^{-\xi t^{-1/Y}(\tilde{U}_t + Z_t)}\right) = e^{\bar{\eta}\xi^Y}, \quad (ii) \lim_{t \rightarrow 0} \tilde{\mathbb{E}}\left(e^{-\xi t^{-1/Y}\tilde{U}_t}\right) = e^{\bar{\eta}^*\xi^Y}, \quad (3.2)$$

where $\bar{\eta} := \Gamma(-Y)(C_+ M^Y + C_- G^Y)$ and $\bar{\eta}^* = \Gamma(-Y)(C_+(M^*)^Y + C_-(G^*)^Y)$. In particular, both $t^{-1/Y}(\tilde{U}_t + Z_t)$ and $t^{-1/Y}\tilde{U}_t$ converge to a Y -stable distribution as $t \rightarrow 0$.

Lemma 3.3. *In the setting and under the assumptions of Section 2, the following two assertions hold true:*

1. *For any $v > 0$,*

$$\begin{aligned} \lim_{t \rightarrow 0} \frac{1}{t} \tilde{\mathbb{P}}\left(Z_t^+ + \tilde{U}_t \geq v\right) &= \int_{\mathbb{R}_0} \mathbf{1}_{\{x - \ln \bar{q}(x) \geq v\}} \tilde{\nu}(dx) \\ &= C_+ \int_0^{\infty} \mathbf{1}_{\{-\ln \bar{q}(x) \geq v\}} x^{-Y-1} dx + C_- \int_{-\infty}^0 \mathbf{1}_{\{-x - \ln \bar{q}(x) \geq v\}} |x|^{-Y-1} dx. \end{aligned} \quad (3.3)$$

2. *There exist constants $\tilde{\kappa} < \infty$ and $t_0 > 0$ such that*

$$(i) \frac{1}{t} \tilde{\mathbb{P}}(\tilde{U}_t \geq v) \leq \tilde{\kappa} v^{-Y}, \quad (ii) \frac{1}{t} \tilde{\mathbb{P}}(Z_t^+ + \tilde{U}_t \geq v) \leq \tilde{\kappa} v^{-Y}, \quad (3.4)$$

for any $0 < t \leq t_0$ and $v > 0$.

We are now in a position to establish the main result of this section. The following result gives the second-order asymptotic behavior of at-the-money call option prices under an exponential tempered stable Lévy model.

Theorem 3.4. *Under the exponential tempered stable model (1.3) without Brownian component ($\sigma = 0$),*

$$\lim_{t \rightarrow 0} t^{\frac{1}{Y}-1} \left(t^{-\frac{1}{Y}} \frac{1}{S_0} \mathbb{E}(S_t - S_0)^+ - \tilde{\mathbb{E}}(Z_1^+) \right) = \vartheta + \eta + \tilde{\gamma} \tilde{\mathbb{P}}(Z_1 \geq 0), \quad (3.5)$$

where η is given as in (2.25), $\tilde{\gamma} := \tilde{\mathbb{E}}L_1$, and

$$\vartheta = C_+ \int_0^{\infty} (1 - \bar{q}(x) + \ln \bar{q}(x)) x^{-Y-1} dx + C_- \int_{-\infty}^0 (1 - e^x \bar{q}(x) + \ln \bar{q}(x) + x) |x|^{-Y-1} dx. \quad (3.6)$$

Remark 3.5. The leading term $\tilde{\mathbb{E}}(Z_1^+)$ can be explicitly computed via the absolute first moment of Z_1 :

$$\tilde{\mathbb{E}}|Z_1| = 2\tilde{\mathbb{E}}(Z_1^+) - \tilde{\mathbb{E}}(Z_1) = 2\tilde{\mathbb{E}}(Z_1^+).$$

Using (1.2.13) in [41], it turns out that

$$\begin{aligned} \tilde{\mathbb{E}}(Z_1^+) &= \frac{1}{2} \tilde{\mathbb{E}}|Z_1| = \frac{1}{\pi} (\hat{C}_+ + \hat{C}_-)^{\frac{1}{Y}} \Gamma\left(1 - \frac{1}{Y}\right) \left(1 + \left(\frac{\hat{C}_+ - \hat{C}_-}{\hat{C}_+ + \hat{C}_-}\right)^2 \tan^2\left(\frac{Y\pi}{2}\right)\right)^{\frac{1}{2Y}} \\ &\quad \times \cos\left(\frac{1}{Y} \arctan\left(\frac{\hat{C}_+ - \hat{C}_-}{\hat{C}_+ + \hat{C}_-} \tan\left(\frac{Y\pi}{2}\right)\right)\right), \end{aligned} \quad (3.7)$$

where $\hat{C}_{\pm} = C_{\pm} \Gamma(-Y) |\cos(\frac{1}{2}Y\pi)|$. In the symmetric case (i.e. $C_- = C_+ = C$), (3.7) takes the form (1.6).

Remark 3.6. From (3.5), it follows that the short-time second-order asymptotic behavior of the ATM call option price (1.2) has the general form:

$$\frac{1}{S_0} \mathbb{E}(S_t - S_0)^+ = d_1 t^{\frac{1}{Y}} + d_2 t + o(t), \quad (t \rightarrow 0), \quad (3.8)$$

with $d_1 = \tilde{\mathbb{E}}(Z_1^+)$ given as in (3.7) and $d_2 = \vartheta + \eta + \tilde{\gamma} \tilde{\mathbb{P}}(Z_1 \geq 0)$. In words, the first-order term synthesizes only the information on the tail-heaviness index Y and the overall jump-intensity parameter, while the second-order term incorporates also the information on the relative intensities of negative and positive jumps as measured by C_+ and C_- .

We now proceed to study the asymptotic behavior of the corresponding Black-Scholes implied volatilities. Throughout, let $\hat{\sigma}(t)$ denote the ATM Black-Scholes implied volatility at maturity t with zero interest rates and dividend yield. The following result gives the asymptotic behavior of $\hat{\sigma}(t)$ as $t \rightarrow 0$.

Proposition 3.7. *Under the exponential tempered stable model (1.3) without Brownian component, the implied volatility $\hat{\sigma}$ has the following small-time behavior:*

$$\hat{\sigma}(t) = \sigma_1 t^{\frac{1}{Y} - \frac{1}{2}} + \sigma_2 t^{\frac{1}{2}} + o(t^{\frac{1}{2}}), \quad t \rightarrow 0, \quad (3.9)$$

where

$$\sigma_1 := \sqrt{2\pi} \tilde{\mathbb{E}}(Z_1^+), \quad (3.10)$$

$$\sigma_2 := \sqrt{2\pi} \left(\vartheta + \eta + \tilde{\gamma} \tilde{\mathbb{P}}(Z_1 \geq 0) \right), \quad (3.11)$$

and where ϑ and η are given by (3.6) and (2.25), respectively.

4 Tempered stable model with Brownian component

In this part, we consider an exponential tempered stable model with non-zero Brownian component. Concretely, throughout, $(X_t)_{t \geq 0}$ is a Lévy process with triplet (σ^2, b, ν) as introduced in Section 2 and $\sigma \neq 0$. In that case, it is well-known that

$$\lim_{t \rightarrow 0} \mathbb{E}^* \left(\exp \left(iu \frac{X_t}{\sqrt{t}} \right) \right) = \exp \left(-\frac{1}{2} \sigma^2 u^2 \right),$$

and, therefore, $X_t/\sqrt{t}$ converges weakly to the centered Gaussian distribution with variance σ^2 . Equivalently, recalling that under $\mathbb{P}^*$, $(W_t^*)_{t \geq 0}$ is a standard Brownian motion, it follows that

$$\lim_{t \rightarrow 0} \mathbb{P}^* \left(\frac{X_t}{\sqrt{t}} \geq x \right) = \mathbb{P}^*(\sigma W_1^* \geq x). \quad (4.1)$$

The first order asymptotic behavior for the ATM European call options in this mixed model is obtained in [43] using Fourier methods. We give next the second-order correction term for the at-the-money European call option price. As before, we change the probability measure $\mathbb{P}^*$ to $\tilde{\mathbb{P}}$ so that $X_t = t\tilde{\gamma} + \sigma W_t^* + Z_t$, with $(Z_t)_{t \geq 0}$ a strictly Y -stable Lévy process under $\tilde{\mathbb{P}}$ (see (2.17)). Recall also that, under both $\mathbb{P}^*$ and $\tilde{\mathbb{P}}$, W^* is still a standard Brownian motion. We will also make use of the decompositions (2.24)-(2.26). The proof of the result below is presented in Appendix B.

Theorem 4.1. *In the setting and under the conditions of Section 2, the ATM European call option price is such that:*

$$\lim_{t \rightarrow 0} t^{\frac{Y}{2} - 1} \left(t^{-\frac{1}{2}} \frac{1}{S_0} \mathbb{E}(S_t - S_0)_+ - \sigma \mathbb{E}^*(W_1^*)_+ \right) = \frac{C_- + C_+}{2Y(Y-1)} \sigma^{1-Y} \mathbb{E}^*(|W_1^*|^{1-Y}). \quad (4.2)$$

Remark 4.2. The $(1-Y)$ -centered moment of a standard normal distribution is given by (see, e.g., (25.6) in [42]):

$$\mathbb{E}^*(|W_1^*|^{1-Y}) = \frac{2^{1-Y}}{\sqrt{\pi}} \Gamma \left(1 - \frac{Y}{2} \right).$$

Thus, the second-order asymptotic behavior of the ATM call option price (1.2) in short-time takes the form

$$\frac{1}{S_0} \mathbb{E}(S_t - S_0)^+ = d_1 t^{\frac{1}{2}} + d_2 t^{\frac{3-Y}{2}} + o\left(t^{\frac{3-Y}{2}}\right), \quad (t \rightarrow 0), \quad (4.3)$$

with

$$d_1 = \frac{\sigma}{\sqrt{2\pi}}, \quad d_2 = \frac{2^{1-Y}}{\sqrt{\pi}} \Gamma\left(1 - \frac{Y}{2}\right) \frac{(C_- + C_+) \sigma^{1-Y}}{2Y(Y-1)}. \quad (4.4)$$

Intuitively, the first-order term only synthesizes the information about the continuous volatility parameter σ , while the second-order term incorporates also the information on the tail index parameter Y and the overall jump-intensity, which is measured by the parameter $\tilde{C} := C_- + C_+$. However, these two-terms do not reflect the individual intensities of negative or positive jumps as measured by the values of C_- and C_+ . This fact suggests that it could be necessary to develop a third-order approximation. For more information on the latter in the CGMY model, we refer the reader to [16].

The next proposition gives the small-time asymptotic behavior for the ATM Black-Scholes implied volatility under the exponential tempered stable model with non-zero Brownian component. Unlike the pure-jump case, we can only derive the first order asymptotics using Theorem 4.1. In fact, the first order term of the ATM call option price under the generalized tempered stable model is the same as the one under the Black-Scholes model. The second order term requires higher order asymptotics of the ATM call option price. The proof is deferred to Appendix B.

Proposition 4.3. *Under the exponential tempered stable model (1.3) with non-zero Brownian component, the implied volatility $\hat{\sigma}$ has the following small-time behavior:*

$$\hat{\sigma}(t) = \sigma + \frac{(C_+ + C_-) 2^{\frac{1}{2}-Y} \sigma^{1-Y}}{Y(Y-1)} \Gamma\left(1 - \frac{Y}{2}\right) t^{1-\frac{Y}{2}} + o\left(t^{1-\frac{Y}{2}}\right), \quad t \rightarrow 0. \quad (4.5)$$

Remark 4.4. A careful analysis of the proofs of the main Theorems 3.4 and 4.1 reveals that the results therein remain true provided that the assertions of Lemmas 3.2 and 3.3 hold true and that the two changes of probability measures of Section 2, first from $\mathbb{P}$ into $\mathbb{P}^*$ and then from $\mathbb{P}^*$ into $\tilde{\mathbb{P}}$, are valid². The first change is valid if (2.1) and (2.4) are satisfied, while, as seen in Remark 2.5, the existence of $\tilde{\mathbb{P}}$ is guaranteed if (2.10) (which in turn implies (2.15)) is satisfied with $M > 1$ and $\ln \tilde{q}^*(x) \leq 0$, for all $x < 0$. In particular, these two conditions are satisfied if (1.11) and

$$q(x) \leq C_+, \quad q(-x) \leq C_-, \quad (x > 0), \quad (4.6)$$

hold true. On the other hand, the proof of Lemma 3.2 will remain valid if, in addition to (1.11) and (4.6), we assume (2.6-iii) and that

$$q(x) \leq C_+ e^{-x}, \quad (x > 0).$$

Finally, Lemma 3.3 does not require any additional assumption.

5 Asymptotics for ATM option prices under a CGMY model

In this section, we specialize the second-order asymptotic expansions of the two previous sections to the CGMY model. The result presented here were first reported in [16]. Recall that under the CGMY model, the Lévy measure of the pure-jump component $(L_t)_{t \geq 0}$ is given by

$$\nu(dx) = |x|^{-Y-1} q(x) dx = |x|^{-Y-1} (C e^{-Mx} \mathbf{1}_{\{x>0\}} + C e^{Gx} \mathbf{1}_{\{x<0\}}) dx, \quad (5.1)$$

with corresponding parameters $C, G, M > 0$ and $Y \in (1, 2)$. Then, the characteristic function of the log-return process $X_t = \sigma W_t + L_t$ takes the form

$$\varphi_t(u) = \mathbb{E}(e^{iuX_t}) = \exp\left(t \left[icu - \frac{\sigma^2 u^2}{2} + C\Gamma(-Y) ((M - iu)^Y + (G + iu)^Y - M^Y - G^Y)\right]\right), \quad (5.2)$$

²Indeed, the first transformation enables us to use the representation of Lemma 3.1, while the second one enables us to exploit the self-similarity of the process $(Z_t)_{t \geq 0}$ and its more explicit distributional behavior in short-time.

for a constant $c \in \mathbb{R}$. The martingale condition (2.1) implies that $M > 1$ and

$$c = -CT(-Y) \left((M-1)^Y + (G+1)^Y - M^Y - G^Y \right) - \frac{\sigma^2}{2}, \quad (5.3)$$

see, e.g., Proposition 4.2 in [43]. In particular, the center $\gamma := \mathbb{E}X_1 = \mathbb{E}L_1$ of X and the parameter b of X (relative to the truncation function $x\mathbf{1}_{\{|x| \leq 1\}}$) are given by

$$\gamma = c - CY\Gamma(-Y)(M^{Y-1} - G^{Y-1}), \quad (5.4)$$

$$b = c - \int_{|x|>1} x\nu(dx) - CY\Gamma(-Y)(M^{Y-1} - G^{Y-1}). \quad (5.5)$$

Under the share measure $\mathbb{P}^*$ introduced in Section 2, $(X_t)_{t \geq 0}$ has Lévy triplet $(b^*, (\sigma^*)^2, \nu^*)$ given by

$$\sigma^* := \sigma, \quad b^* := c + \sigma^2 - \int_{|x|>1} x\nu^*(dx) - CY\Gamma(-Y)((M^*)^{Y-1} - (G^*)^{Y-1}), \quad \nu^*(dx) := e^x \nu(dx), \quad (5.6)$$

with $M^* = M-1$ and $G^* = G+1$. Finally, under the probability measure $\tilde{\mathbb{P}}$, the centered process $(Z_t)_{t \geq 0}$ is symmetric, and its center $\tilde{\gamma} := \tilde{\mathbb{E}}X_1 = \tilde{\mathbb{E}}L_1^*$ is given by (see [16] for the detailed computation)

$$\begin{aligned} \tilde{\gamma} &= \tilde{b} + \int_{\{|x|>1\}} x\tilde{\nu}(dx) = b^* + \int_{|x| \leq 1} x(\tilde{\nu} - \nu^*)(dx) + \int_{\{|x|>1\}} x\tilde{\nu}(dx) \\ &= -CT(-Y) \left((M-1)^Y + (G+1)^Y - M^Y - G^Y \right) + \frac{\sigma^2}{2}. \end{aligned} \quad (5.7)$$

We also need the value of η defined in (2.25), which, under the CGMY model, is now given by

$$\begin{aligned} \eta &= C \int_{0+}^{\infty} (e^{-M^*x} - 1 + M^*x)x^{-Y-1}dx + C \int_{-\infty}^{0-} (e^{G^*x} - 1 - G^*x)|x|^{-Y-1}dx \\ &= CT(-Y) \left((M^*)^Y + (G^*)^Y \right), \end{aligned} \quad (5.8)$$

where, for the last equality above, we used the analytic continuation of (14.19) in [42].

We are ready to explicitly write the second-order expansions. First, let us compute the term ϑ of (3.6). To this end, it is convenient to use the representation given in (A.15) below noting that $\bar{q}(x) := e^{-Mx}\mathbf{1}_{\{x>0\}} + e^{Gx}\mathbf{1}_{\{x<0\}}$ in the CGMY case:

$$\begin{aligned} \vartheta &= C \int_0^{\infty} (e^{-v} - 1) \int_{\mathbb{R}_0} \mathbf{1}_{\{x - \ln \bar{q}(x) \geq v\}} |x|^{-Y-1} dx dv \\ &= C \int_0^{\infty} (e^{-v} - 1) \int_0^{\infty} \mathbf{1}_{\{Mx \geq v\}} x^{-Y-1} dx dv + C \int_0^{\infty} (e^{-v} - 1) \int_{-\infty}^0 \mathbf{1}_{\{-x - Gx \geq v\}} (-x)^{-Y-1} dx dv \\ &= \frac{C}{Y} \left(M^Y + (G+1)^Y \right) \int_0^{\infty} (e^{-v} - 1) v^{-Y} dv \\ &= -CT(-Y) \left((M^*+1)^Y + (G^*)^Y \right), \end{aligned} \quad (5.9)$$

where in the last equality we used the following identity (see p. 84 in [42]):

$$\int_0^{\infty} (e^{-v} - 1) v^{-Y} dv = \Gamma(1-Y) = -Y\Gamma(-Y).$$

Using (5.7)-(5.9), it follows that in the pure-jump CGMY model,

$$d_2 := \vartheta + \eta + \frac{\tilde{\gamma}}{2} = \frac{CT(-Y)}{2} \left((M-1)^Y - M^Y - (G+1)^Y + G^Y \right),$$

while in the general CGMY model with non-zero Brownian component, $C_+ = C_- = C$, and thus

$$d_2 := \frac{C\sigma^{1-Y}2^{1-Y}}{Y(Y-1)\sqrt{\pi}} \Gamma\left(1 - \frac{Y}{2}\right)$$

6 Numerical examples

In this part, we assess the performance of the previous approximations through a detailed numerical analysis for a CGMY model.

6.1 The numerical methods

Let us first select a suitable numerical method to compute the ATM option prices by considering two methods: Inverse Fourier Transform (IFT) and Monte Carlo (MC).

Before introducing the IFT method, let us set some notations. The characteristic function corresponding to the Black-Scholes model with volatility Σ is given by

$$\varphi_t^{BS,\Sigma}(v) = \exp\left(-\frac{\Sigma^2 t}{2}(v^2 + iv)\right).$$

The corresponding call option price at the log-moneyness $k = \log(S_0/K)$ under the Black-Scholes model with volatility Σ is denoted by $C_{BS}^\Sigma(k)$; that is,

$$C_{BS}^\Sigma(k) = S_0 e^{-rt} \mathbb{E}(e^{(r-\Sigma^2/2)t + \Sigma W_t} - e^k)_+.$$

Let us also recall that the characteristic function under the generalized CGMY model is denoted by φ_t (see (5.2)) and let us denote the corresponding call option price at log-moneyness k by $C(k)$. The IFT method is based on the following inversion formula (see Section 11.1.3 in [8]):

$$z_T(k) := C(k) - C_{BS}^\Sigma(k) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-ivk} \zeta_T(v) dv, \quad (6.1)$$

where

$$\zeta_T(v) := e^{ivr} \frac{\varphi_T(v-i) - \varphi_T^{BS,\Sigma}(v-i)}{iv(1+iv)}. \quad (6.2)$$

In our case, we fix $r = 0$ and, since we are only interested in ATM option prices, we set $k = 0$. In order to compute numerically the integral in (6.1), we use Simpson's rule:

$$z_T(0) := \frac{1}{2\pi} \int_{-\infty}^{\infty} \zeta_T(v) dv = \Delta \sum_{m=0}^{P-1} w_m^* \zeta_T(v_m^*),$$

with $\Delta = Q/(P-1)$, $v_m^* = -Q/2 + m\Delta$, and $w_0 = 1/2$, $w_{2\ell-1} = 4/3$, and $w_{2\ell} = 2/3$, for $\ell = 1, \dots, P/2$.

We also consider a Monte Carlo method based on the risk-neutral option price representation under the probability measure $\tilde{\mathbb{P}}$. Under this probability measure and using the notation (2.25) as well as the relations (2.24) and (2.26), we have:

$$\begin{aligned} \mathbb{E}(e^{X_T} - 1)_+ &= \mathbb{E}^* \left(e^{-X_T} (e^{X_T} - 1)_+ \right) = \tilde{\mathbb{E}} \left(e^{-U_T} (1 - e^{-X_T})_+ \right) \\ &= \tilde{\mathbb{E}} \left(e^{-M^* \bar{U}_T^+ + G^* \bar{U}_T^- - \eta T} \left(1 - e^{-\bar{U}_T^+ - \bar{U}_T^- - T\tilde{\gamma} - \sigma W_T} \right)_+ \right), \end{aligned}$$

which can be easily computed by Monte Carlo method using the fact that, under $\tilde{\mathbb{P}}$, the variables $\bar{U}_T^+$ and $-\bar{U}_T^-$ are independent Y -stable random variables with scale, skewness, and location parameters $TC|\cos(\pi Y/2)|\Gamma(-Y)$, 1, and 0, respectively. Standard simulation methods are available to generate stable random variables. We consider the following set of parameters

$$C = 0.5, \quad G = 2, \quad M = 3.6, \quad Y = 1.5.$$

Figure 1 compares the first- and second-order approximations as given in Remarks 3.6 and 4.2 to the prices based on the Inverse Fourier Transform (IFT-based price) and the Monte Carlo method (MC-based price) under both the pure-jump case and the generalized CGMY case with $\sigma = 0.4$. For the MC-based price, we use 100,000 simulations, while for the IFT-based method, we use $P = 2^{14}$ and $Q = 800$. As it can be seen, it is not easy to integrate numerically the characteristic function (6.2) since T is quite small and, therefore, the characteristic functions φ_T and $\varphi_T^{BS,\Sigma}$ are quite flat. The Monte Carlo method turns out to be much more accurate and faster.




Figure 1: Comparisons of ATM call option prices for two methods (Inverse Fourier Transform and Monte-Carlo method) with the first- and second-order approximations. MC-based price is based on 100,000 simulations while the IFT-based method is based on the parameter values $P = 2^{14}$ and $Q = 800$. The parameter σ in the generalized CGMY model is set to be 0.1.

6.2 Results for different parameter settings

Here, we investigate the performance of the approximations for different settings of parameters:

1. Figure 2 compares the 1st- and 2nd-order approximations with the MC prices for different values of C , fixing the values of all the other parameters. In the pure-jump case, the 2nd order approximation is significantly better for moderately small values of C , but for larger values of C , this is not the case unless T is extremely small. For a nonzero continuous component, the 1st order approximation is extremely bad as it only takes into account the parameter σ .
2. Figure 3 compares the 1st- and 2nd-order approximations with the MC prices for different values of Y , fixing the values of all the other parameters. In both cases, the 2nd order approximation is significantly better for values of Y around 1.5, which is consistent with the observation that $|d_2| \rightarrow \infty$ as $Y \rightarrow 1$ or $Y \rightarrow 2$. For a nonzero continuous component, the 1st order approximation is again extremely bad as compared to the 2nd order approximation.
3. In the left panel of Figure 4, we analyze the effect of the relative intensities of negative jumps compared to positive jumps in the pure-jump CGMY case. That is, we fix the values M to be 4 and consider different values for G . As expected, since the first order approximation does not take into account this information, the 2nd-order approximation performs significantly better.
4. In the right panel of Figure 4, we analyze the effect of the volatility of the continuous component in the generalized CGMY case. The 2nd order approximation is, in general, much better than the 1st-order approximation and, interestingly enough, the quality of the 2nd order approximations improves as the values of σ increases. In fact, it seems that the 2nd-order approximation and the MC prices collapse to a steady curve as σ increases.




Figure 2: Comparisons of ATM call option prices with the short-time approximations for different values of the jump intensity parameter C .




Figure 3: Comparisons of ATM call option prices with the short-time approximations for different values of the tail-heaviness parameter Y .




Figure 4: Comparisons of ATM call option prices with the short-time approximations for different values of G or M and different values of the volatility parameter σ .

A Proofs of Section 3: Pure-jump tempered stable model

For simplicity, unless otherwise stated, throughout this section, we fix $S_0 = 1$.

Proof of Lemma 3.1.

From (1.9), we have

$$\begin{aligned} t^{-1/Y} \mathbb{E}(S_t - S_0)^+ &= t^{-1/Y} \mathbb{P}^*(X_t \geq E) = t^{-1/Y} \int_0^\infty e^{-x} \mathbb{P}^*(X_t \geq x) dx \\ &= \int_0^\infty e^{-t^{1/Y} u} \mathbb{P}^*(t^{-1/Y} X_t \geq u) du. \end{aligned} \quad (\text{A.1})$$

Next, using the change of probability measure (2.23),

$$\mathbb{P}^*(t^{-1/Y} X_t \geq u) = \mathbb{E}^*(\mathbf{1}_{\{t^{-1/Y} X_t \geq u\}}) = \tilde{\mathbb{E}}(e^{-U_t} \mathbf{1}_{\{t^{-1/Y} X_t \geq u\}}),$$

and, moreover, from (2.17) and (2.24),

$$\mathbb{P}^*(t^{-1/Y} X_t \geq u) = e^{-\eta t} \tilde{\mathbb{E}}(e^{-\tilde{U}_t} \mathbf{1}_{\{t^{-1/Y} Z_t \geq u - \tilde{\gamma} t^{1-1/Y}\}}).$$

Then,

$$t^{-1/Y} \mathbb{E}(S_t - S_0)^+ = \int_0^\infty e^{-t^{1/Y} u - \eta t} \tilde{\mathbb{E}}(e^{-\tilde{U}_t} \mathbf{1}_{\{t^{-1/Y} Z_t \geq u - \tilde{\gamma} t^{1-1/Y}\}}) du, \quad (\text{A.2})$$

and, changing variables ($v = u - \tilde{\gamma} t^{1-1/Y}$), the result follows. $\square$

Proof of Lemma 3.2.

Fix $Q_t := \tilde{U}_t + Z_t$. From the usual moment generating function formula for Poisson integrals, the change of variables $y = t^{-1/Y}x$, and recalling that $\varphi(x) + x = -\ln \bar{q}(x)$,

$$\begin{aligned}\tilde{\mathbb{E}}\left(e^{-\xi t^{-1/Y}Q_t}\right) &= \tilde{\mathbb{E}}\left(\exp\left\{-\xi t^{-1/Y}\int_0^t\int(\varphi(x)+x)\bar{N}(ds,dx)\right\}\right) \\ &= \exp\left\{t\int\left(e^{\xi t^{-1/Y}\ln\bar{q}(x)}-1-\xi t^{-1/Y}\ln\bar{q}(x)\right)\tilde{\nu}(dx)\right\} \\ &= \exp\left\{C_+\int_0^\infty\left(e^{\xi t^{-1/Y}\ln\bar{q}(t^{1/Y}y)}-1-\xi t^{-1/Y}\ln\bar{q}(t^{1/Y}y)\right)y^{-Y-1}dy\right. \\ &\quad \left.+C_-\int_{-\infty}^0\left(e^{\xi t^{-1/Y}\ln\bar{q}(t^{1/Y}y)}-1-\xi t^{-1/Y}\ln\bar{q}(t^{1/Y}y)\right)|y|^{-Y-1}dy\right\}.\end{aligned}\quad (\text{A.3})$$

Let $\Xi(u) := e^u - 1 - u$ and $B := \sup|\ln\bar{q}(w)|/|w|$, which is necessarily finite in light of (2.6-iii) and (2.11-ii). Now, using that $\ln\bar{q}(u) \leq 0$, $u \in \mathbb{R}_0$, and that $|\Xi(u)| \leq K(u^2 \wedge |u|)$, for $u \leq 0$ and some constant $K < \infty$,

$$\begin{aligned}\Xi\left(\xi t^{-\frac{1}{Y}}\ln\bar{q}\left(t^{\frac{1}{Y}}y\right)\right)|y|^{-Y-1} &\leq K\left(\left(\xi t^{-\frac{1}{Y}}\ln\bar{q}\left(t^{\frac{1}{Y}}y\right)\right)^2 \wedge \left|\xi t^{-\frac{1}{Y}}\ln\bar{q}\left(t^{\frac{1}{Y}}y\right)\right|\right)|y|^{-Y-1} \\ &\leq K\left((\xi y B)^2 \wedge |\xi y B|\right)|y|^{-Y-1} \\ &\leq K\left((\xi B)^2 \vee |\xi B|\right)(y^2 \wedge |y|)|y|^{-Y-1},\end{aligned}$$

which is integrable. Therefore, one can pass the limit inside the integrals in (A.3) and, using (2.11),

$$\begin{aligned}\lim_{t \rightarrow 0}\tilde{\mathbb{E}}\left(e^{-\xi t^{-1/Y}Q_t}\right) &= \exp\left\{C_+\int_0^\infty(e^{-y\xi M}-1+y\xi M)y^{-Y-1}dy+C_-\int_{-\infty}^0(e^{-|y|\xi G}-1+|y|\xi G)|y|^{-Y-1}dy\right\} \\ &= \exp\left\{\left(C_+\int_0^\infty(e^{-uM}-1+uM)u^{-Y-1}du+C_-\int_{-\infty}^0(e^{-|u|G}-1+|u|G)|u|^{-Y-1}du\right)\xi^Y\right\}.\end{aligned}$$

Finally, using the analytic continuation of the representation (14.19) in [42], we show that the last expression is of the form $\exp(\eta^*\xi^Y)$, with η^* given as in the statement of the Lemma.

For (3.2-ii), we proceed as above to get

$$\begin{aligned}\tilde{\mathbb{E}}\left(e^{-\xi t^{-1/Y}\tilde{U}_t}\right) &= \exp\left\{C_+\int_0^\infty\Xi\left(\xi t^{-1/Y}\ln\bar{q}^*\left(t^{1/Y}y\right)\right)y^{-Y-1}dy\right. \\ &\quad \left.+C_-\int_{-\infty}^0\Xi\left(\xi t^{-1/Y}\ln\bar{q}^*\left(t^{1/Y}y\right)\right)|y|^{-Y-1}dy\right\}.\end{aligned}\quad (\text{A.4})$$

Now, for all $y \in \mathbb{R}_0$,

$$\xi t^{-1/Y}\ln\bar{q}^*\left(t^{1/Y}y\right) = \xi t^{-1/Y}\ln\bar{q}\left(t^{1/Y}y\right) + \xi y = \xi y\left(\frac{1}{yt^{1/Y}}\ln\bar{q}\left(t^{1/Y}y\right) + 1\right) \leq 0. \quad (\text{A.5})$$

Indeed, this is clear when $y < 0$, since $\ln\bar{q}(u) \leq 0$, while for $y > 0$, (A.5) follows from

$$\sup_{x>0}\frac{\ln\bar{q}(x)}{x} \leq -1,$$

since, from (2.5), for all $x > 0$,

$$e^x\bar{q}(x) = e^x\int_0^1e^{-x/s}s^Y\bar{\rho}(ds) = \int_0^1e^{-x(\frac{1}{s}-1)}s^Y\bar{\rho}(ds) \leq \int_0^1s^Y\bar{\rho}(ds) = 1.$$

Using (A.5), we can proceed as above to justify passing the limit into the integrals in (A.4). Next, using Lemma 2.4,

$$\begin{aligned}\lim_{t \rightarrow 0}\tilde{\mathbb{E}}\left(e^{-\xi t^{-1/Y}\tilde{U}_t}\right) &= \exp\left\{C_+\int_0^\infty(e^{-y\xi M^*}-1+y\xi M^*)y^{-Y-1}dy+C_-\int_{-\infty}^0(e^{-|y|\xi G^*}-1+|y|\xi G^*)|y|^{-Y-1}dy\right\} \\ &= \exp\left\{\left(C_+\int_0^\infty(e^{-uM^*}-1+uM^*)u^{-Y-1}du+C_-\int_{-\infty}^0(e^{-|u|G^*}-1+|u|G^*)|u|^{-Y-1}du\right)\xi^Y\right\}.\end{aligned}$$

Finally, (3.2-ii) follows once more from the analytic continuation of (14.19) in [42]. $\square$

Proof of Lemma 3.3.

We use a small/large jump type decomposition. Concretely, fix $\varepsilon > 0$ and let

$$\bar{Z}_t^{(\varepsilon)} := \int_0^t \int_{|x| \geq \varepsilon} x N(ds, dx), \quad Z_t^{(\varepsilon)} := Z_t - \bar{Z}_t^{(\varepsilon)}.$$

Denote respectively by $(N_t^{(\varepsilon)})_{t \geq 0}$ and $(\xi_i^{(\varepsilon)})_{i \geq 1}$ the counting process and sizes of the jumps of $(\bar{Z}_t^{(\varepsilon)})_{t \geq 0}$, so that $\bar{Z}_t^{(\varepsilon)} = \sum_{i=1}^{N_t^{(\varepsilon)}} \xi_i^{(\varepsilon)}$. Next, defined the corresponding processes for $\tilde{U}$:

$$\tilde{U}_t^{(\varepsilon)} := \int_0^t \int_{|x| \geq \varepsilon} \varphi(x) N(ds, dx) = - \sum_{i=1}^{N_t^{(\varepsilon)}} \ln \bar{q}^* \left(\xi_i^{(\varepsilon)} \right), \quad \tilde{U}_t^{(\varepsilon)} := \tilde{U}_t - \tilde{U}_t^{(\varepsilon)}.$$

Let $\lambda_\varepsilon := \mathbb{E} N_1^{(\varepsilon)}$ be the jumps intensity of $\bar{Z}^{(\varepsilon)}$. We prove the validity of the two assertions in two steps:

(1) By conditioning on $N_t^{(\varepsilon)}$,

$$\begin{aligned} t^{-1} \tilde{\mathbb{P}} \left(Z_t^+ + \tilde{U}_t \geq v \right) &= t^{-1} \tilde{\mathbb{P}} \left(\left(Z_t^{(\varepsilon)} \right)^+ + \tilde{U}_t^{(\varepsilon)} \geq v \right) e^{-\lambda_\varepsilon t} \\ &\quad + e^{-\lambda_\varepsilon t} \lambda_\varepsilon \tilde{\mathbb{P}} \left(\left(Z_t^{(\varepsilon)} + \xi_1^{(\varepsilon)} \right)^+ + \tilde{U}_t^{(\varepsilon)} - \ln \bar{q}^* \left(\xi_1^{(\varepsilon)} \right) \geq v \right) + O(t). \end{aligned}$$

The first term above can be made $O(t)$ by taking $0 < \varepsilon < \varepsilon_0$, for some $\varepsilon_0 > 0$ (see, e.g., [38, Lemma 3.2] or [42, Section 26]). For the second term, first note that for some $\varepsilon_0 > 0$, small enough,

$$\begin{aligned} a_0(v) &:= \lambda_\varepsilon \tilde{\mathbb{P}} \left(\left(\xi_1^{(\varepsilon)} \right)^+ - \ln \bar{q}^* \left(\xi_1^{(\varepsilon)} \right) \geq v \right) \\ &= \int_{\mathbb{R}_0} \mathbf{1}_{\{x^+ - \ln \bar{q}^*(x) \geq v\}} \tilde{\nu}(dx) = \int_{\mathbb{R}_0} \mathbf{1}_{\{x^- - \ln \bar{q}(x) \geq v\}} \tilde{\nu}(dx) \\ &= C_+ \int_0^\infty \mathbf{1}_{\{-\ln \bar{q}(x) \geq v\}} x^{-Y-1} (dx) + C_- \int_{-\infty}^0 \mathbf{1}_{\{-x - \ln \bar{q}(x) \geq v\}} |x|^{-Y-1} dx, \end{aligned}$$

for all $0 < \varepsilon < \varepsilon_0$. Also, since

$$F_{v,\varepsilon}(z, u) := \tilde{\mathbb{P}} \left(\left(z + \xi_1^{(\varepsilon)} \right)^+ + u - \ln \bar{q}^* \left(\xi_1^{(\varepsilon)} \right) \geq v \right)$$

is continuous at $(z, u) = (0, 0)$, for any fixed $0 < \varepsilon < \varepsilon_0$, the function

$$A_t(v) := t^{-1} \tilde{\mathbb{P}} \left(Z_t^+ + \tilde{U}_t \geq v \right) - a_0(v)$$

is such that

$$\begin{aligned} \lim_{t \rightarrow 0} A_t(v) &= \lambda_\varepsilon \lim_{t \rightarrow 0} \left(\tilde{\mathbb{P}} \left(\left(Z_t^{(\varepsilon)} + \xi_1^{(\varepsilon)} \right)^+ + \tilde{U}_t^{(\varepsilon)} - \ln \bar{q}^* \left(\xi_1^{(\varepsilon)} \right) \geq v \right) - \lambda_\varepsilon \tilde{\mathbb{P}} \left(\left(\xi_1^{(\varepsilon)} \right)^+ - \ln \bar{q}^* \left(\xi_1^{(\varepsilon)} \right) \geq v \right) \right) \\ &= \lambda_\varepsilon \lim_{t \rightarrow 0} \tilde{\mathbb{E}} \left(F_{v,\varepsilon} \left(Z_t^{(\varepsilon)}, \tilde{U}_t^{(\varepsilon)} \right) - F_{v,\varepsilon}(0, 0) \right) \\ &= \lambda_\varepsilon \tilde{\mathbb{E}} \left(\lim_{t \rightarrow 0} F_{v,\varepsilon} \left(Z_t^{(\varepsilon)}, \tilde{U}_t^{(\varepsilon)} \right) - F_{v,\varepsilon}(0, 0) \right) \\ &= 0, \end{aligned}$$

where we had used the dominated convergence theorem to obtain the last equality.

(2) Throughout, $\tilde{\kappa} > 0$ denotes a generic constant that may vary from line to line. First, note that (2.22) implies that

$$\frac{1}{t} \tilde{\mathbb{P}}(Z_t^+ \geq v) \leq \tilde{\kappa} v^{-Y}, \quad (\text{A.6})$$

for any $0 < t \leq 1$ and $v > 0$ and some constants $\tilde{\kappa} < \infty$. So, it suffices to show the analog inequality for $\tilde{U}$.

Let $\bar{C} := C_+ + C_-$. Using the decompositions in the previous step (1) with $\varepsilon = v/4$,

$$\frac{1}{t} \tilde{\mathbb{P}}\left(\tilde{U}_t^{(\varepsilon)} \geq v/2\right) \leq \frac{1}{t} \tilde{\mathbb{P}}\left(N_t^{(\varepsilon)} \neq 0\right) = \frac{1}{t} (1 - e^{-\lambda_\varepsilon t}) \leq \lambda_\varepsilon \leq \bar{C} \int_{|x| \geq v/4} |x|^{-Y-1} dx \leq \tilde{\kappa} v^{-Y},$$

for some $\tilde{\kappa} < \infty$. Then,

$$\tilde{\mathbb{E}} \tilde{U}_t^{(\varepsilon)} = t \int_{|x| > \varepsilon} \ln \bar{q}^*(x) \tilde{\nu}(dx) = t \int_{|x| > \varepsilon} \ln \bar{q}(x) \tilde{\nu}(dx) + t C_- \int_{x < -\varepsilon} x |x|^{-Y-1} dx + t C_+ \int_{x > \varepsilon} x |x|^{-Y-1} dx,$$

and, since $\bar{q}(x) \leq 1$, for all x , $\tilde{\mathbb{E}} \tilde{U}_t^{(\varepsilon)} \leq t C_+ \int_{x > \varepsilon} x |x|^{-Y-1} dx = C_+(Y-1)^{-1} 4^{Y-1} t v^{1-Y}$. Thus, whenever t and v are such that $C_+(Y-1)^{-1} 4^{Y-1} t v^{1-Y} \leq v/4$, we have

$$\tilde{\mathbb{P}}(\tilde{U}_t^{(\varepsilon)} \geq v/2) \leq \tilde{\mathbb{P}}(\tilde{U}_t^{(\varepsilon)} - \tilde{\mathbb{E}} \tilde{U}_t^{(\varepsilon)} + \tilde{\mathbb{E}} \tilde{U}_t^{(\varepsilon)} \geq v/2) \leq \tilde{\mathbb{P}}(\tilde{U}_t^{(\varepsilon)} - \tilde{\mathbb{E}} \tilde{U}_t^{(\varepsilon)} \geq v/4).$$

Next, using a concentration inequality for centered random variables (see, e.g., [22], Corollary 1),

$$\tilde{\mathbb{P}}(\tilde{U}_t^{(\varepsilon)} \geq v/2) \leq \tilde{\mathbb{P}}(\tilde{U}_t^{(\varepsilon)} - \tilde{\mathbb{E}} \tilde{U}_t^{(\varepsilon)} \geq v/4) \leq e^{\frac{v}{4\varepsilon} - \left(\frac{v}{4\varepsilon} + \frac{tV_\varepsilon^2}{\varepsilon^2}\right) \log\left(1 + \frac{\varepsilon v}{4tV_\varepsilon^2}\right)} \leq \left(\frac{4eV_\varepsilon^2}{\varepsilon v}\right)^{\frac{v}{4\varepsilon}} t^{\frac{v}{4\varepsilon}} \leq \frac{eV_{v/4}^2}{v^2} t,$$

where $V_\varepsilon^2 := \text{Var}(\tilde{U}_t^{(\varepsilon)})$ and in the last inequality we recall that $\varepsilon = v/4$. Now,

$$V_\varepsilon^2 = \text{Var}(\tilde{U}_t^{(\varepsilon)}) = \int_{\{|x| \leq \varepsilon\}} (\ln \bar{q}^*(x))^2 \tilde{\nu}(dx) \leq C \int_{\{|x| \leq \varepsilon\}} x^2 |x|^{-Y-1} dx \leq \tilde{\kappa} v^{2-Y},$$

for some constants $\tilde{\kappa} < \infty$, where above, we set $C = \sup_{|x| > 0} |\ln \bar{q}^*(x)|/|x|$, which is finite in light of (2.6-iii) and (2.15-ii). Therefore, whenever $C_+(Y-1)^{-1} 4^{Y-1} t v^{1-Y} \leq v/4$ (or equivalently, $C_+(Y-1)^{-1} 4^Y t v^{-Y} \leq 1$),

$$\frac{1}{t} \tilde{\mathbb{P}}(\tilde{U}_t^{(\varepsilon)} \geq v/2) \leq \frac{eV_{v/4}^2}{v^2} \leq \tilde{\kappa} v^{-Y}.$$

Moreover, for any $t > 0$ and $v > 0$,

$$\begin{aligned} \frac{1}{t} \tilde{\mathbb{P}}(\tilde{U}_t^{(\varepsilon)} \geq v/2) &= \frac{1}{t} \tilde{\mathbb{P}}(\tilde{U}_t^{(\varepsilon)} \geq v/2) \mathbf{1}_{\{C_+(Y-1)^{-1} 4^Y t v^{-Y} \leq 1\}} + \frac{1}{t} \tilde{\mathbb{P}}(\tilde{U}_t^{(\varepsilon)} \geq v/2) \mathbf{1}_{\{C_+(Y-1)^{-1} 4^Y t v^{-Y} > 1\}} \\ &\leq \tilde{\kappa} v^{-Y} + \frac{C_+}{t} (Y-1)^{-1} 4^Y t v^{-Y} \leq \tilde{\kappa}' v^{-Y}, \end{aligned}$$

for some constant $\tilde{\kappa}'$. Combining the previous estimates, we finally have

$$\frac{1}{t} \tilde{\mathbb{P}}(\tilde{U}_t \geq v) \leq \frac{1}{t} \tilde{\mathbb{P}}(\tilde{U}_t^{(\varepsilon)} \geq v/2) + \frac{1}{t} \tilde{\mathbb{P}}(\tilde{U}_t^{(\varepsilon)} \geq v/2) \leq \tilde{\kappa} v^{-Y}, \quad (\text{A.7})$$

for all $v > 0$ and $t > 0$ and some constant $\tilde{\kappa} < \infty$. □

Proof of Theorem 3.4.

For simplicity, we first treat the case $\tilde{\gamma} = 0$ so that, in light of Lemma 3.1,

$$t^{-1/Y} \mathbb{E}(S_t - S_0)^+ = e^{-\eta t} \int_0^\infty e^{-t^{1/Y} v} \tilde{\mathbb{E}}\left(e^{-\tilde{U}_t} \mathbf{1}_{\{t^{-1/Y} Z_t \geq v\}}\right) dv. \quad (\text{A.8})$$

The general case is resolved in Lemma A.1 below. Let

$$D(t) := t^{-1/Y} \mathbb{E}(S_t - S_0)^+ - \tilde{\mathbb{E}}(Z_1^+),$$

which can be written as

$$\begin{aligned} D(t) &:= \int_0^\infty e^{-t^{1/Y}v} \tilde{\mathbb{E}}\left(e^{-\tilde{U}_t} \mathbf{1}_{\{t^{-1/Y}Z_t \geq v\}}\right) dv - \tilde{\mathbb{E}}(Z_1^+) \\ &\quad + (e^{-\eta t} - 1) \tilde{\mathbb{E}}(Z_1^+) \\ &\quad + (e^{-\eta t} - 1) \left(\int_0^\infty e^{-t^{1/Y}v} \tilde{\mathbb{E}}\left(e^{-\tilde{U}_t} \mathbf{1}_{\{t^{-1/Y}Z_t \geq v\}}\right) dv - \tilde{\mathbb{E}}(Z_1^+) \right) \\ &=: D_1(t) + D_2(t) + D_3(t). \end{aligned} \tag{A.9}$$

We will show that

$$t^{1/Y-1} D_1(t) \longrightarrow \vartheta + \eta, \quad \text{as } t \rightarrow 0, \tag{A.10}$$

for a certain constant ϑ , while it is clear that $D_3(t) = o(D_1(t))$ and that $t^{1/Y-1} D_2(t) = o(1)$, as $t \rightarrow 0$. First,

$$\begin{aligned} D_1(t) &= \tilde{\mathbb{E}}\left(e^{-\tilde{U}_t} \int_0^{t^{-1/Y}Z_t^+} e^{-t^{1/Y}v} dv\right) - \tilde{\mathbb{E}}(Z_1^+) \\ &= t^{-1/Y} \left(\tilde{\mathbb{E}}\left(e^{-\tilde{U}_t}\right) - \tilde{\mathbb{E}}\left(e^{-(\tilde{U}_t + Z_t^+)}\right) \right) - \tilde{\mathbb{E}}(Z_1^+) \\ &= t^{-1/Y} \left(e^{\eta t} - \tilde{\mathbb{E}}\left(e^{-(\tilde{U}_t + Z_t^+)}\right) \right) - \tilde{\mathbb{E}}(Z_1^+). \end{aligned} \tag{A.11}$$

Next, by the self-similarity of $(Z_t)_{t \geq 0}$ under $\tilde{\mathbb{P}}$ (see (2.18)), $\tilde{\mathbb{E}}(Z_1^+) = t^{-1/Y} \tilde{\mathbb{E}}(Z_t^+)$ and, also using that $\tilde{\mathbb{E}}(\tilde{U}_t) = 0$,

$$\begin{aligned} t^{1/Y-1} D_1(t) &= t^{1/Y-1} \left(\frac{e^{\eta t} - 1}{t^{1/Y}} + \frac{1 - \tilde{\mathbb{E}}\left(e^{-(Z_t^+ + \tilde{U}_t)}\right) - \tilde{\mathbb{E}}\left(Z_t^+ + \tilde{U}_t\right)}{t^{1/Y}} \right) \\ &= \frac{e^{\eta t} - 1}{t} + \frac{1}{t} \tilde{\mathbb{E}}\left(\int_0^{Z_t^+ + \tilde{U}_t} (e^{-v} - 1) dv \mathbf{1}_{\{Z_t^+ + \tilde{U}_t \geq 0\}}\right) - \frac{1}{t} \tilde{\mathbb{E}}\left(\int_{Z_t^+ + \tilde{U}_t}^0 (e^{-v} - 1) dv \mathbf{1}_{\{Z_t^+ + \tilde{U}_t \leq 0\}}\right) \\ &= \frac{e^{\eta t} - 1}{t} + \frac{1}{t} \int_0^\infty (e^{-v} - 1) \tilde{\mathbb{P}}(Z_t^+ + \tilde{U}_t \geq v) dv - \frac{1}{t} \int_0^\infty (e^v - 1) \tilde{\mathbb{P}}(Z_t^+ + \tilde{U}_t \leq -v) dv \\ &=: D_{11}(t) + D_{12}(t) + D_{13}(t). \end{aligned} \tag{A.12}$$

Clearly,

$$D_{11}(t) \rightarrow \eta, \quad \text{as } t \rightarrow 0. \tag{A.13}$$

Let $Q_t := Z_t + \tilde{U}_t$ and note that $Q_t \leq Z_t^+ + \tilde{U}_t$. Then, for $D_{13}(t)$, using that $e^y - 1 \leq ye^y$, $y > 0$, Markov's inequality, and since $1 < Y < 2$,

$$\begin{aligned} 0 \leq D_{13}(t) &\leq t^{1/Y-1} \int_0^\infty \left(e^{t^{1/Y}u} - 1\right) \tilde{\mathbb{P}}\left(t^{-1/Y}Q_t \leq -u\right) du \\ &\leq t^{2/Y-1} \int_0^\infty e^{t^{1/Y}u} u \tilde{\mathbb{P}}\left(t^{-1/Y}Q_t \leq -u\right) du \\ &\leq t^{2/Y-1} \int_0^\infty e^{(t^{1/Y}-1)u} u du \tilde{\mathbb{E}}\left(e^{-t^{-1/Y}Q_t}\right) \xrightarrow{t \rightarrow 0} 0, \end{aligned} \tag{A.14}$$

where in the last step we applied (3.2-i).

Let us now deal with D_{12} . First, from (3.4-ii), one can apply dominated convergence and pass the limit inside the integrals so that

$$\lim_{t \rightarrow 0} D_{12}(t) = \int_0^\infty (e^{-v} - 1) \lim_{t \rightarrow 0} \left(t^{-1} \tilde{\mathbb{P}}(Z_t^+ + \tilde{U}_t \geq v) \right) dv.$$

From (3.3), we then have

$$\lim_{t \rightarrow 0} D_{12}(t) = \int_0^\infty (e^{-v} - 1) \int_{\mathbb{R}_0} \mathbf{1}_{\{x^- - \ln \bar{q}(x) \geq v\}} \tilde{\nu}(dx) dv := \vartheta. \quad (\text{A.15})$$

Combining (A.12), (A.13), (A.14), and (A.15), it follows that

$$\lim_{t \rightarrow 0} t^{1/Y-1} D_1(t) = \int_0^\infty (e^{-v} - 1) \int_{\mathbb{R}_0} \mathbf{1}_{\{x^- - \ln \bar{q}(x) \geq v\}} \tilde{\nu}(dx) dv + \eta.$$

Finally, the expression in (3.6) follows recalling that $\tilde{\nu}(dx) = |x|^{-Y-1} (C_+ \mathbf{1}_{\{x>0\}} + C_- \mathbf{1}_{\{x<0\}}) dx$ and applying Fubini's theorem to the right-hand side of (A.15):

$$\begin{aligned} & C_+ \int_0^\infty (e^{-v} - 1) \int_0^\infty \mathbf{1}_{\{-\ln \bar{q}(x) \geq v\}} x^{-Y-1} dx dv + C_- \int_0^\infty (e^{-v} - 1) \int_{-\infty}^0 \mathbf{1}_{\{-x - \ln \bar{q}(x) \geq v\}} |x|^{-Y-1} dx dv \\ &= C_+ \int_0^\infty \int_0^{-\ln \bar{q}(x)} (e^{-v} - 1) dv x^{-Y-1} dx + C_- \int_{-\infty}^0 \int_0^{-x - \ln \bar{q}(x)} (e^{-v} - 1) dv |x|^{-Y-1} dx \\ &= C_+ \int_0^\infty \left(1 - e^{\ln \bar{q}(x)} + \ln \bar{q}(x) \right) x^{-Y-1} dx + C_- \int_{-\infty}^0 \left(1 - e^{x + \ln \bar{q}(x)} + x + \ln \bar{q}(x) \right) |x|^{-Y-1} dx. \end{aligned}$$

This concludes the proof. $\square$

Lemma A.1. *If $\tilde{\gamma} \neq 0$ in (3.1), then*

$$\lim_{t \rightarrow 0} t^{\frac{1}{Y}-1} \left(t^{-\frac{1}{Y}} \frac{1}{S_0} \mathbb{E}(S_t - S_0)^+ - S_0 \tilde{\mathbb{E}}(Z_1^+) \right) = \vartheta + \eta + \tilde{\gamma} \tilde{\mathbb{P}}(Z_1 \geq 0). \quad (\text{A.16})$$

Proof. Without loss of generality, fix $S_0 = 1$ and also assume that $\tilde{\gamma} > 0$ (the case $\tilde{\gamma} < 0$ being similar). Using (3.1),

$$\begin{aligned} \lim_{t \rightarrow 0} t^{1/Y-1} \left(t^{-1/Y} \mathbb{E}(S_t - S_0)^+ - \tilde{\mathbb{E}}(Z_1^+) \right) &= \lim_{t \rightarrow 0} t^{1/Y-1} \left(e^{-(\tilde{\gamma}+\eta)t} \int_0^\infty e^{-t^{1/Y} v} \tilde{\mathbb{E}} \left(e^{-\tilde{U}_t} \mathbf{1}_{\{t^{-1/Y} Z_t \geq v\}} \right) dv - \tilde{\mathbb{E}}(Z_1^+) \right) \\ &\quad + \lim_{t \rightarrow 0} t^{1/Y-1} e^{-(\tilde{\gamma}+\eta)t} \int_{-\tilde{\gamma} t^{1-1/Y}}^0 e^{-t^{1/Y} v} \tilde{\mathbb{E}} \left(e^{-\tilde{U}_t} \mathbf{1}_{\{t^{-1/Y} Z_t \geq v\}} \right) dv \\ &=: \tilde{D}_{11}(t) + \tilde{D}_{12}(t). \end{aligned}$$

As in the proof of (A.10),

$$\lim_{t \rightarrow 0} \tilde{D}_{11}(t) = \vartheta + \eta. \quad (\text{A.17})$$

For $\tilde{D}_{12}(t)$, changing variables to $u = t^{1/Y-1}$ and probability measure to $\mathbb{P}^*$, we have

$$\tilde{D}_{12}(t) = e^{-\tilde{\gamma}t} \int_{-\tilde{\gamma}}^0 e^{-tu} \tilde{\mathbb{E}} \left(e^{-(\tilde{U}_t + \eta t)} \mathbf{1}_{\{Z_t \geq tu\}} \right) du = e^{-\tilde{\gamma}t} \int_{-\tilde{\gamma}}^0 e^{-tu} \mathbb{P}^* \left(t^{-1/Y} Z_t \geq t^{1-1/Y} u \right) du.$$

Next, recall that $(L_t^*)_{t \geq 0}$ and, therefore, also $(Z_t)_{t \geq 0}$, are tempered stable Lévy processes under $\mathbb{P}^*$, as shown by (2.13)-(2.14). Then, since $Y \in (1, 2)$, $t^{-1/Y} Z_t$ converges in distribution to a Y -stable random variable $\tilde{Z}$ under $\mathbb{P}^*$ with center (or mean) 0 and spectral measure

$$\sigma(+1) = \int_0^\infty u^Y \rho(du) = \int_0^\infty s^Y \rho(ds) = C_+, \quad \sigma(-1) = \int_{-\infty}^0 |u|^Y \rho(du) = \int_{-\infty}^0 |s|^Y \rho(ds) = C_-,$$

see Theorem 3.1 in [39] (see also Proposition 1 in [37]). Hence, the distribution of $\tilde{Z}$ (under $\mathbb{P}^*$) is the same as the distribution of Z_1 under $\tilde{\mathbb{P}}$. Thus, Slutsky's lemma implies that $t^{-1/Y}Z_t - t^{1-1/Y}u \xrightarrow{\mathcal{D}} \tilde{Z}$ and, in particular,

$$\lim_{t \rightarrow 0} \mathbb{P}^* \left(t^{-1/Y}Z_t - t^{1-1/Y}u \geq 0 \right) = \mathbb{P}^* \left(\tilde{Z} \geq 0 \right) = \tilde{\mathbb{P}}(Z_1 \geq 0).$$

Finally, by the dominated convergence theorem,

$$\lim_{t \rightarrow 0} \tilde{D}_{12}(t) = \tilde{\gamma} \tilde{\mathbb{P}}(Z_1 \geq 0). \quad (\text{A.18})$$

Combining (A.18) with (A.17) leads to (A.16). $\square$

Proof of Proposition 3.7.

The small-time asymptotic behavior of the ATM call option price $C_{BS}(t, \sigma)$ at maturity t under the Black-Scholes model with volatility σ and zero interest rates is (recall that $S_0 = 1$)

$$C_{BS}(t, \sigma) = \frac{\sigma}{\sqrt{2\pi}} t^{1/2} - \frac{\sigma^3}{24\sqrt{2\pi}} t^{3/2} + O(t^{5/2}), \quad t \rightarrow 0; \quad (\text{A.19})$$

see, e.g., [19, Corollary 3.4]. To derive the small-time asymptotics for the implied volatility, we need a result analogous to (A.19) when σ is replaced by $\hat{\sigma}(t)$. The following representation taken from [36, Lemma 3.1] will be useful,

$$C_{BS}(t, \sigma) = F(\sigma\sqrt{t}) \quad \text{with} \quad F(\theta) := \int_0^\theta \Phi' \left(\frac{v}{2} \right) dv = \frac{1}{\sqrt{2\pi}} \int_0^\theta \exp \left(-\frac{v^2}{8} \right) dv,$$

together with the Taylor expansion for F at $\theta = 0$ (see [36, Lemma 5.1]); i.e.,

$$F(\theta) = \frac{1}{\sqrt{2\pi}} \theta - \frac{1}{24\sqrt{2\pi}} \theta^3 + O(\theta^5), \quad \theta \rightarrow 0.$$

Then, since $\hat{\sigma}(t) \rightarrow 0$ as $t \rightarrow 0$ (see, e.g., [43, Proposition 5]), we conclude that

$$C_{BS}(t, \hat{\sigma}(t)) = \frac{\hat{\sigma}(t)}{\sqrt{2\pi}} t^{1/2} - \frac{\hat{\sigma}(t)^3}{24\sqrt{2\pi}} t^{3/2} + O \left((\hat{\sigma}(t) t^{1/2})^5 \right), \quad \text{as } t \rightarrow 0. \quad (\text{A.20})$$

Returning to the proof of Proposition 3.7, by equating (3.8) and (A.20) and comparing the first order terms,

$$\tilde{\mathbb{E}}(Z_1^+) t^{1/Y} \sim \frac{\hat{\sigma}(t)}{\sqrt{2\pi}} t^{1/2}, \quad t \rightarrow 0,$$

and, therefore,

$$\hat{\sigma}(t) \sim \sqrt{2\pi} \tilde{\mathbb{E}}(Z_1^+) t^{\frac{1}{Y} - \frac{1}{2}} := \sigma_1 t^{\frac{1}{Y} - \frac{1}{2}}, \quad t \rightarrow 0. \quad (\text{A.21})$$

Next, set $\tilde{\sigma}(t) = \hat{\sigma}(t) - \sigma_1 t^{\frac{1}{Y} - \frac{1}{2}}$. By comparing the first and second order terms in (3.8) with the first term in (A.20) (noting that the second order term in (A.20) is $o(t)$),

$$\left(\vartheta + \eta + \tilde{\gamma} \tilde{\mathbb{P}}(Z_1 \geq 0) \right) t \sim \frac{\tilde{\sigma}(t)}{\sqrt{2\pi}} t^{1/2}, \quad t \rightarrow 0.$$

Hence, $\tilde{\sigma}(t) \rightarrow 0$ as $t \rightarrow 0$, and moreover

$$\tilde{\sigma}(t) \sim \sqrt{2\pi} \left(\vartheta + \eta + \tilde{\gamma} \tilde{\mathbb{P}}(Z_1 \geq 0) \right) t^{1/2}, \quad t \rightarrow 0. \quad (\text{A.22})$$

Combining (A.21) and (A.22) finishes the proof. $\square$

B Proofs of Section 4: Tempered stable with Brownian component

Proof of Theorem 4.1.

For simplicity, fix $S_0 = 1$. Recalling that $X_t = \sigma W_t^* + L_t^*$ under $\mathbb{P}^*$, and using (1.9), the self-similarity of W^* , and the change of variable $u = t^{-1/2}x$,

$$R_t := t^{-1/2} \mathbb{E}(S_t - S_0)_+ - \sigma \mathbb{E}^*(W_1^*)_+ = \int_0^\infty e^{-\sqrt{t}u} \mathbb{P}^*(\sigma W_1^* \geq u - t^{-\frac{1}{2}} L_t^*) du - \int_0^\infty \mathbb{P}^*(\sigma W_1^* \geq u) du.$$

Next, changing the probability measure to $\tilde{\mathbb{P}}$, using that $L_t^* = Z_t + \tilde{\gamma}t$, $U_t = \tilde{U}_t + \eta t$, and the change of variable $y = u - t^{1/2}\tilde{\gamma}$ in the first integral above, lead to

$$\begin{aligned} R_t &= \int_{-t^{1/2}\tilde{\gamma}}^\infty e^{-\sqrt{t}y - \tilde{\gamma}t} \tilde{\mathbb{E}} e^{-\tilde{U}_t - \eta t} \mathbf{1}_{\{\sigma W_1^* \geq y - t^{-\frac{1}{2}} Z_t\}} dy - \int_0^\infty \tilde{\mathbb{E}} e^{-\tilde{U}_t - \eta t} \mathbf{1}_{\{\sigma W_1^* \geq u\}} du \\ &= e^{-(\eta + \tilde{\gamma})t} \int_0^\infty e^{-\sqrt{t}y} \left(\tilde{\mathbb{E}} e^{-\tilde{U}_t} \mathbf{1}_{\{\sigma W_1^* \geq y - t^{-\frac{1}{2}} Z_t\}} - \tilde{\mathbb{E}} e^{-\tilde{U}_t} \mathbf{1}_{\{\sigma W_1^* \geq y\}} \right) dy \\ &\quad + e^{-(\eta + \tilde{\gamma})t} \int_{-t^{1/2}\tilde{\gamma}}^0 e^{-\sqrt{t}y} \tilde{\mathbb{E}} e^{-\tilde{U}_t} \mathbf{1}_{\{\sigma W_1^* \geq y - t^{-\frac{1}{2}} Z_t\}} dy \\ &\quad + \int_0^\infty \left(e^{-\tilde{\gamma}t - \sqrt{t}y} - 1 \right) \mathbb{P}^*(\sigma W_1^* \geq y) dy. \end{aligned} \tag{B.1}$$

Above, the last term is clearly $O(t^{1/2})$ as $t \rightarrow 0$, while the second term can be shown to be asymptotically equivalent to a term that is $O(t^{1/2})$ by arguments analogous to those of (A.18). Thus, we only need to study the term in (B.1) that we denote A_t and which can be written as:

$$A_t = e^{-\tilde{\eta}t} \tilde{\mathbb{E}} \left(e^{-\tilde{U}_t} \int_0^\infty e^{-\sqrt{t}y} \left(\mathbf{1}_{\{\sigma W_1^* \geq y - t^{-\frac{1}{2}} Z_t\}} - \mathbf{1}_{\{\sigma W_1^* \geq y\}} \right) dy \right),$$

where we had set $\tilde{\eta} := \eta + \tilde{\gamma}$. To study the asymptotic behavior of A_t , decompose it into the following three parts

$$\begin{aligned} A_t &= e^{-\tilde{\eta}t} \tilde{\mathbb{E}} \left(e^{-\tilde{U}_t} \mathbf{1}_{\{W_1^* \geq 0, \sigma W_1^* + t^{-\frac{1}{2}} Z_t \geq 0\}} \int_{\sigma W_1^*}^{\sigma W_1^* + t^{-\frac{1}{2}} Z_t} e^{-\sqrt{t}y} dy \right) \\ &\quad - e^{-\tilde{\eta}t} \tilde{\mathbb{E}} \left(e^{-\tilde{U}_t} \mathbf{1}_{\{0 \leq \sigma W_1^* \leq -t^{-\frac{1}{2}} Z_t\}} \int_0^{\sigma W_1^*} e^{-\sqrt{t}y} dy \right) \\ &\quad + e^{-\tilde{\eta}t} \tilde{\mathbb{E}} \left(e^{-\tilde{U}_t} \mathbf{1}_{\{0 \leq -\sigma W_1^* \leq t^{-\frac{1}{2}} Z_t\}} \int_0^{\sigma W_1^* + t^{-\frac{1}{2}} Z_t} e^{-\sqrt{t}y} dy \right) \\ &=: I_1(t) - I_2(t) + I_3(t). \end{aligned} \tag{B.2}$$

We analyze each of there three terms in the following three steps:

Step 1. We first study the behavior of $I_1(t)$. Since $(Z_t)_{t \geq 0}$ and $(W_t^*)_{t \geq 0}$ are independent,

$$\begin{aligned} I_1(t) &= e^{-\tilde{\eta}t} \tilde{\mathbb{E}} \left(\mathbf{1}_{\{W_1^* \geq 0, \sigma W_1^* + t^{-\frac{1}{2}} Z_t \geq 0\}} \frac{e^{-\tilde{U}_t} - e^{-(\tilde{U}_t + Z_t)}}{\sqrt{t}} e^{-\sqrt{t}\sigma W_1^*} \right) \\ &= e^{-\tilde{\eta}t} \int_0^\infty \tilde{\mathbb{E}} \left(\mathbf{1}_{\{Z_t \geq -t^{\frac{1}{2}}y\}} \frac{e^{-\tilde{U}_t} (1 - e^{-Z_t})}{\sqrt{t}} \right) e^{-\sqrt{t}y} \frac{e^{-\frac{y^2}{2\sigma^2}}}{\sqrt{2\pi\sigma^2}} dy \\ &=: e^{-\tilde{\eta}t} \int_0^\infty J_1(t, y) e^{-\sqrt{t}y} \frac{e^{-\frac{y^2}{2\sigma^2}}}{\sqrt{2\pi\sigma^2}} dy \end{aligned} \tag{B.3}$$

Using the self-similarity of $(Z_t)_{t \geq 0}$ and since $\tilde{\mathbb{E}}Z_t = 0$, $J_1(t, y)$ is then decomposed as:

$$\begin{aligned} J_1(t, y) &= \tilde{\mathbb{E}} \left(\mathbf{1}_{\{Z_t \geq -t^{\frac{1}{2}}y\}} \left(\frac{e^{-\tilde{U}_t} - e^{-(\tilde{U}_t + Z_t)}}{\sqrt{t}} - t^{-\frac{1}{2}}Z_t \right) \right) + t^{\frac{1}{Y}-\frac{1}{2}}\tilde{\mathbb{E}} \left((-Z_1) \mathbf{1}_{\{-Z_1 \geq t^{\frac{1}{2}-\frac{1}{Y}}y\}} \right) \\ &=: J_{11}(t, y) + J_{12}(t, y). \end{aligned} \quad (\text{B.4})$$

Let us first consider $J_{12}(t, y)$. From (2.19)-(2.21), there exists a constant $\lambda > 0$ such that

$$t^{\frac{Y}{2}-1}J_{12}(t, y) \leq \lambda y^{1-Y}, \quad (\text{B.5})$$

for any $0 < t \leq 1$ and $y > 0$ (see Appendix C for the verification of this claim). Moreover, for any fixed $y > 0$,

$$t^{\frac{Y}{2}-1}J_{12}(t, y) = t^{\frac{Y}{2}+\frac{1}{Y}-\frac{3}{2}} \int_{t^{\frac{1}{2}-\frac{1}{Y}}y}^{\infty} up_Z(-u)du = t^{\frac{Y}{2}-\frac{1}{Y}-\frac{1}{2}} \int_y^{\infty} wp_Z(-t^{\frac{1}{2}-\frac{1}{Y}}w)dw.$$

Using (2.19), there exists $0 < t_0 < 1$ such that

$$t^{\frac{Y}{2}-\frac{1}{Y}-\frac{1}{2}}wp_Z(-t^{\frac{1}{2}-\frac{1}{Y}}w) \leq 2(C_+ \vee C_-)w^{-Y},$$

for any $0 < t < t_0$ and $w \geq y$. Therefore, by the dominated convergence theorem, and in light of (2.19), we get:

$$\begin{aligned} &\lim_{t \rightarrow 0} t^{\frac{Y}{2}-1} e^{-\tilde{\eta}t} \int_0^{\infty} J_{12}(t, y) e^{-\sqrt{t}y} \frac{e^{-\frac{y^2}{2\sigma^2}}}{\sqrt{2\pi\sigma^2}} dy \\ &= \int_0^{\infty} \left(\lim_{t \rightarrow 0} t^{\frac{Y}{2}-1} J_{12}(t, y) \right) \frac{e^{-\frac{y^2}{2\sigma^2}}}{\sqrt{2\pi\sigma^2}} dy \\ &= \int_0^{\infty} \left(\int_y^{\infty} w \left(\lim_{t \rightarrow 0} t^{\frac{Y}{2}-\frac{1}{Y}-\frac{1}{2}} p_Z(-t^{\frac{1}{2}-\frac{1}{Y}}w) \right) dw \right) \frac{e^{-\frac{y^2}{2\sigma^2}}}{\sqrt{2\pi\sigma^2}} dy \\ &= C_- \int_0^{\infty} \left(\int_y^{\infty} w^{-Y} dw \right) \frac{e^{-\frac{y^2}{2\sigma^2}}}{\sqrt{2\pi\sigma^2}} dy = \frac{C_-}{Y-1} \int_0^{\infty} y^{1-Y} \frac{e^{-\frac{y^2}{2\sigma^2}}}{\sqrt{2\pi\sigma^2}} dy. \end{aligned} \quad (\text{B.6})$$

For $J_{11}(t, y)$,

$$\begin{aligned} J_{11}(t, y) &= t^{-\frac{1}{2}}\tilde{\mathbb{E}} \left(\mathbf{1}_{\{Z_t \geq 0\}} \int_{\tilde{U}_t}^{\tilde{U}_t + Z_t} (e^{-x} - 1) dx \right) - t^{-\frac{1}{2}}\tilde{\mathbb{E}} \left(\mathbf{1}_{\{-t^{\frac{1}{2}}y \leq Z_t \leq 0\}} \int_{\tilde{U}_t + Z_t}^{\tilde{U}_t} (e^{-x} - 1) dx \right) \\ &= t^{-\frac{1}{2}} \int_{\mathbb{R}} (e^{-x} - 1) T_1(t, x, y) dx - t^{-\frac{1}{2}} \int_{\mathbb{R}} (e^{-x} - 1) T_2(t, x, y) dx, \end{aligned} \quad (\text{B.7})$$

where, for $t > 0$ and $y > 0$, we set

$$T_1(t, x, y) := \tilde{\mathbb{P}}(Z_t \geq 0, \tilde{U}_t \leq x \leq \tilde{U}_t + Z_t), \quad T_2(t, x, y) := \tilde{\mathbb{P}}(-t^{\frac{1}{2}}y \leq Z_t \leq 0, \tilde{U}_t + Z_t \leq x \leq \tilde{U}_t).$$

By (3.4-ii), there exists $\tilde{\kappa} > 0$, such that for any $x > 0$ and $0 < t \leq 1$,

$$T_1(t, x, y) \leq \tilde{\mathbb{P}}(x \leq \tilde{U}_t + Z_t) \leq \tilde{\mathbb{P}}(x \leq \tilde{U}_t + Z_t^+) \leq \tilde{\kappa} t x^{-Y}. \quad (\text{B.8})$$

Hence,

$$\begin{aligned} 0 &\leq e^{-\tilde{\eta}t} t^{\frac{Y}{2}-1} \int_0^{\infty} \int_0^{\infty} \frac{(1 - e^{-x})}{\sqrt{t}} T_1(t, x, y) dx \frac{e^{-\sqrt{t}y} e^{-\frac{y^2}{2\sigma^2}}}{\sqrt{2\pi\sigma^2}} dy \\ &\leq e^{-\tilde{\eta}t} \tilde{\kappa} t^{\frac{Y-1}{2}} \int_0^{\infty} \int_0^{\infty} (1 - e^{-x}) x^{-Y} dx \frac{e^{-\frac{y^2}{2\sigma^2}}}{\sqrt{2\pi\sigma^2}} dy \longrightarrow 0, \end{aligned} \quad (\text{B.9})$$

as $t \rightarrow 0$, since $Y > 1$. Similarly, using (3.4-i), there exists a constant $\tilde{\kappa} < \infty$ such that

$$T_2(t, x, y) \leq \tilde{\mathbb{P}}(\tilde{U}_t \geq x) \leq \tilde{\kappa} t x^{-Y}, \quad (\text{B.10})$$

for any $x > 0$ and $0 < t \leq 1$, and, thus, as in (B.9),

$$\lim_{t \rightarrow 0} \frac{e^{-\tilde{\eta}t}}{t^{1-\frac{Y}{2}}} \int_0^\infty \int_0^\infty \frac{(1-e^{-x})}{\sqrt{t}} T_2(t, x, y) dx \frac{e^{-\sqrt{t}y} e^{-\frac{y^2}{2\sigma^2}}}{\sqrt{2\pi\sigma^2}} dy = 0. \quad (\text{B.11})$$

For $x < 0$, using (3.2-ii) and Markov's inequality, there exist $0 < t_0 < 1$ and $\tilde{\kappa} < \infty$ such that

$$T_1(t, x, y) \leq \tilde{\mathbb{P}}(t^{-1/Y} \tilde{U}_t \leq t^{-\frac{1}{Y}} x) \leq \tilde{\mathbb{E}}(e^{-t^{-1/Y} \tilde{U}_t}) e^{t^{-1/Y} x} \leq \tilde{\kappa} e^{t^{-1/Y} x}, \quad (\text{B.12})$$

for any $0 < t \leq t_0$. Therefore,

$$0 \leq t^{\frac{Y-3}{2}} \int_{-\infty}^0 (e^{-x} - 1) T_1(t, x, y) dx \leq \tilde{\kappa} t^{\frac{Y-3}{2}} \int_{-\infty}^0 \left(e^{(t^{-\frac{1}{Y}} - 1)x} - e^{t^{-\frac{1}{Y}} x} \right) dx = \tilde{\kappa} t^{\frac{Y-3}{2}} \frac{t^{\frac{2}{Y}}}{1 - t^{\frac{1}{Y}}}.$$

Hence, by the dominated convergence theorem,

$$0 \leq e^{-\tilde{\eta}t} t^{\frac{Y}{2}-1} \int_0^\infty \int_{-\infty}^0 \frac{e^{-x} - 1}{\sqrt{t}} T_1(t, x, y) dx \frac{e^{-\sqrt{t}y} e^{-\frac{y^2}{2\sigma^2}}}{\sqrt{2\pi\sigma^2}} dy \leq \tilde{\kappa} t^{\frac{Y-3}{2}} \frac{t^{\frac{2}{Y}}}{1 - t^{\frac{1}{Y}}} \int_0^\infty \frac{e^{-\frac{y^2}{2\sigma^2}}}{\sqrt{2\pi\sigma^2}} dy \rightarrow 0, \quad (\text{B.13})$$

as $t \rightarrow 0$, since $2/Y > 1 > (3-Y)/2$, for $1 < Y < 2$. Similarly, using (3.2-i), for $x < 0$,

$$T_2(t, x, y) \leq \tilde{\mathbb{P}}(\tilde{U}_t + Z_t \leq x) \leq \tilde{\mathbb{E}}\left(e^{-t^{-1/Y}(\tilde{U}_t + Z_t)}\right) e^{t^{-1/Y} x} \leq \tilde{\kappa} e^{t^{-1/Y} x},$$

for any $0 < t \leq t_0$ and some constant $\tilde{\kappa}$. Therefore, as in (B.13),

$$\lim_{t \rightarrow 0} e^{-\tilde{\eta}t} t^{\frac{Y}{2}-1} \int_0^\infty \int_{-\infty}^0 \frac{(1-e^{-x})}{\sqrt{t}} T_2(t, x, y) dx \frac{e^{-\sqrt{t}y} e^{-\frac{y^2}{2\sigma^2}}}{\sqrt{2\pi\sigma^2}} dy = 0. \quad (\text{B.14})$$

Combining (B.6), (B.9), (B.11), (B.13) and (B.14), we finally obtain

$$\lim_{t \rightarrow 0} t^{\frac{Y}{2}-1} I_1(t) = \frac{C_-}{Y-1} \int_0^\infty y^{1-Y} \frac{e^{-\frac{y^2}{2\sigma^2}}}{\sqrt{2\pi\sigma^2}} dy. \quad (\text{B.15})$$

Step 2. Next, we study the asymptotic behavior of $I_2(t)$. Using the independence of $(Z_t)_{t \geq 0}$ and $(W_t^*)_{t \geq 0}$,

$$\begin{aligned} I_2(t) &= e^{-\tilde{\eta}t} \tilde{\mathbb{E}} \left(e^{-\tilde{U}_t} \mathbf{1}_{\{0 \leq \sigma W_1^* \leq -t^{-\frac{1}{2}} Z_t\}} \frac{1 - e^{-\sqrt{t}\sigma W_1^*}}{\sqrt{t}} \right) \\ &= e^{-\tilde{\eta}t} \int_0^\infty \tilde{\mathbb{E}} \left(e^{-\tilde{U}_t} \mathbf{1}_{\{Z_t \leq -t^{\frac{1}{2}} y\}} \right) \frac{1 - e^{-\sqrt{t}y}}{\sqrt{t}} \frac{e^{-\frac{y^2}{2\sigma^2}}}{\sqrt{2\pi\sigma^2}} dy \\ &= e^{-\tilde{\eta}t} \int_0^\infty \tilde{\mathbb{E}} \left((e^{-\tilde{U}_t} - 1) \mathbf{1}_{\{Z_t \leq -t^{\frac{1}{2}} y\}} \right) \frac{1 - e^{-\sqrt{t}y}}{\sqrt{t}} \frac{e^{-\frac{y^2}{2\sigma^2}}}{\sqrt{2\pi\sigma^2}} dy \\ &\quad + e^{-\tilde{\eta}t} \int_0^\infty \tilde{\mathbb{P}} \left(Z_t \leq -t^{\frac{1}{2}} y \right) \frac{1 - e^{-\sqrt{t}y}}{\sqrt{t}} \frac{e^{-\frac{y^2}{2\sigma^2}}}{\sqrt{2\pi\sigma^2}} dy. \end{aligned} \quad (\text{B.16})$$

By (2.22) and the self-similarity of $(Z_t)_{t \geq 0}$, for $y > 0$,

$$\tilde{\mathbb{P}} \left(Z_t \leq -t^{\frac{1}{2}} y \right) \frac{1 - e^{-\sqrt{t}y}}{\sqrt{t}} = \tilde{\mathbb{P}} \left(Z_1 \leq -t^{\frac{1}{2}-\frac{1}{Y}} y \right) \frac{1 - e^{-\sqrt{t}y}}{\sqrt{t}} \leq \kappa t^{1-\frac{Y}{2}} y^{1-Y} \leq \kappa y^{1-Y},$$

which, when multiplied by $\exp(-y^2/2\sigma^2)$, becomes integrable on $[0, \infty)$. Hence, by (2.21) and the dominated convergence theorem,

$$\lim_{t \rightarrow 0} t^{\frac{Y}{2}-1} e^{-\tilde{\eta}t} \int_0^\infty \tilde{\mathbb{P}}\left(Z_t \leq -t^{\frac{1}{2}}y\right) \frac{1 - e^{-\sqrt{t}y}}{\sqrt{t}} \frac{e^{-\frac{y^2}{2\sigma^2}}}{\sqrt{2\pi\sigma^2}} dy = \frac{C_-}{Y} \int_0^\infty y^{1-Y} \frac{e^{-\frac{y^2}{2\sigma^2}}}{\sqrt{2\pi\sigma^2}} dy. \quad (\text{B.17})$$

To find the asymptotic behavior of the first integral in (B.16), we decompose it as:

$$\begin{aligned} \tilde{\mathbb{E}}\left((e^{-\tilde{U}_t} - 1)\mathbf{1}_{\{Z_t \leq -t^{\frac{1}{2}}y\}}\right) &= \tilde{\mathbb{E}}\left(\mathbf{1}_{\{Z_t \leq -t^{\frac{1}{2}}y, \tilde{U}_t < 0\}} \int_{\tilde{U}_t}^0 e^{-u} du\right) - \tilde{\mathbb{E}}\left(\mathbf{1}_{\{Z_t \leq -t^{\frac{1}{2}}y, \tilde{U}_t \geq 0\}} \int_0^{\tilde{U}_t} e^{-u} du\right) \\ &=: J_{21}(t, y) + J_{22}(t, y). \end{aligned} \quad (\text{B.18})$$

For $J_{21}(t, y)$, using Markov's inequality and (3.2-ii), there exist $\tilde{\kappa} < \infty$ and $0 < t_0 < 1$ such that

$$\begin{aligned} J_{21}(t, y) &= \int_{-\infty}^0 e^{-x} \tilde{\mathbb{P}}\left(Z_t \leq -t^{\frac{1}{2}}y, t^{-\frac{1}{Y}}\tilde{U}_t \leq t^{-\frac{1}{Y}}x\right) dx \\ &\leq \tilde{\mathbb{E}}(e^{-t^{-1/Y}\tilde{U}_t}) \int_{-\infty}^0 e^{-x} \exp\left(t^{-\frac{1}{Y}}x\right) dx \leq \tilde{\kappa} \frac{t^{\frac{1}{Y}}}{1 - t^{\frac{1}{Y}}}, \end{aligned}$$

for any $0 < t < t_0$ and $y \geq 0$. Since $1 - Y/2 < 1/2 < 1/Y$, for $1 < Y < 2$, by the dominated convergence theorem,

$$0 \leq t^{\frac{Y}{2}-1} e^{-\tilde{\eta}t} \int_0^\infty J_{21}(t, y) \frac{1 - e^{-\sqrt{t}y}}{\sqrt{t}} \frac{e^{-\frac{y^2}{2\sigma^2}}}{\sqrt{2\pi\sigma^2}} dy \leq \tilde{\kappa} t^{\frac{Y}{2}-1} \frac{t^{\frac{1}{Y}}}{1 - t^{\frac{1}{Y}}} \int_0^\infty y \frac{e^{-\frac{y^2}{2\sigma^2}}}{\sqrt{2\pi\sigma^2}} dy \longrightarrow 0. \quad (\text{B.19})$$

We further decompose the second term $J_{22}(t, y)$ in (B.18) as:

$$\begin{aligned} J_{22}(t, y) &= \tilde{\mathbb{E}}\left((e^{-\tilde{U}_t} - 1 + \tilde{U}_t)\mathbf{1}_{\{Z_t \leq -t^{\frac{1}{2}}y, \tilde{U}_t \geq 0\}}\right) - \tilde{\mathbb{E}}\left(\tilde{U}_t \mathbf{1}_{\{Z_t \leq -t^{\frac{1}{2}}y, \tilde{U}_t \geq 0\}}\right) \\ &=: J_{22}^{(1)}(t, y) - J_{22}^{(2)}(t, y). \end{aligned} \quad (\text{B.20})$$

Using (3.4-i) and the fact that $1 - Y/2 < 1/2 < 1/Y$, for $1 < Y < 2$,

$$\begin{aligned} 0 \leq J_{22}^{(2)}(t, y) &\leq t^{\frac{1}{Y}} \tilde{\mathbb{E}}\left(t^{-1/Y}\tilde{U}_t\right)^+ \\ &= t^{1/Y} \int_0^\infty \mathbb{P}\left(t^{-1/Y}\tilde{U}_t \geq u\right) du \end{aligned} \quad (\text{B.21})$$

$$\leq t^{1/Y} \left(1 + \int_1^\infty \mathbb{P}\left(t^{-1/Y}\tilde{U}_t \geq u\right) du\right) \quad (\text{B.22})$$

$$\leq t^{1/Y} \left(1 + \int_1^\infty \tilde{\kappa} t(t^{1/Y}u)^{-Y} du\right) \quad (\text{B.23})$$

$$\leq K t^{1/Y}, \quad (\text{B.24})$$

which is clearly $o(t^{1-\frac{Y}{2}})$. Moreover,

$$\begin{aligned} J_{22}^{(1)}(t, y) &= \tilde{\mathbb{E}}\left(\int_0^{\tilde{U}_t} (1 - e^{-w}) dw \mathbf{1}_{\{Z_t \leq -t^{\frac{1}{2}}y, \tilde{U}_t \geq 0\}}\right) \\ &= \int_0^\infty (1 - e^{-w}) \tilde{\mathbb{P}}\left(\tilde{U}_t \geq w, Z_t \leq -t^{\frac{1}{2}}y\right) dw \\ &\leq \int_0^\infty (1 - e^{-w}) \tilde{\mathbb{P}}\left(\tilde{U}_t \geq w\right) dw. \end{aligned}$$

Using (B.10), $(1 - e^{-\sqrt{t}y})/\sqrt{t} \leq y$, for $y > 0$, and applying the dominated convergence theorem, we get

$$0 \leq t^{\frac{Y}{2}-1} e^{-\tilde{\eta}t} \int_0^\infty J_{22}^{(1)}(t, y) \frac{1 - e^{-\sqrt{t}y}}{\sqrt{t}} \frac{e^{-\frac{y^2}{2\sigma^2}}}{\sqrt{2\pi\sigma^2}} dy \leq t^{\frac{Y}{2}} \int_0^\infty (1 - e^{-w}) w^{-Y} dw \int_0^\infty y \frac{e^{-\frac{y^2}{2\sigma^2}}}{\sqrt{2\pi\sigma^2}} dy \longrightarrow 0, \quad (\text{B.25})$$

as $t \rightarrow 0$. Combining (B.17), (B.19), (B.24) and (B.25), lead to

$$\lim_{t \rightarrow 0} t^{\frac{Y}{2}-1} I_2(t) = \frac{C_-}{Y} \int_0^\infty y^{1-Y} \frac{e^{-\frac{y^2}{2\sigma^2}}}{\sqrt{2\pi\sigma^2}} dy. \quad (\text{B.26})$$

Step 3. We finally study the behavior of $I_3(t)$. Note that

$$\begin{aligned} I_3(t) &= e^{-\tilde{\eta}t} \int_0^\infty \tilde{\mathbb{E}} \left(e^{-\tilde{U}_t} \mathbf{1}_{\{Z_t \geq t^{\frac{1}{2}}y\}} \frac{1 - e^{\sqrt{t}y} e^{-Z_t}}{\sqrt{t}} \right) \frac{e^{-\frac{y^2}{2\sigma^2}}}{\sqrt{2\pi\sigma^2}} dy \\ &= e^{-\tilde{\eta}t} \int_0^\infty \tilde{\mathbb{E}} \left(e^{-\tilde{U}_t} \mathbf{1}_{\{Z_t \geq t^{\frac{1}{2}}y\}} \right) \frac{1 - e^{\sqrt{t}y}}{\sqrt{t}} \frac{e^{-\frac{y^2}{2\sigma^2}}}{\sqrt{2\pi\sigma^2}} dy \\ &\quad + e^{-\tilde{\eta}t} \int_0^\infty \tilde{\mathbb{E}} \left(e^{-\tilde{U}_t} \mathbf{1}_{\{Z_t \geq t^{\frac{1}{2}}y\}} \frac{1 - e^{-Z_t}}{\sqrt{t}} \right) e^{\sqrt{t}y} \frac{e^{-\frac{y^2}{2\sigma^2}}}{\sqrt{2\pi\sigma^2}} dy. \end{aligned} \quad (\text{B.27})$$

We first investigate $J_{31}(t, y) := \tilde{\mathbb{E}} \left(e^{-\tilde{U}_t} \mathbf{1}_{\{Z_t \geq t^{\frac{1}{2}}y\}} \right)$ by decomposing it as:

$$\begin{aligned} J_{31}(t, y) &= \tilde{\mathbb{E}} \left((e^{-\tilde{U}_t} - 1) \mathbf{1}_{\{Z_t \geq t^{\frac{1}{2}}y\}} \right) + \tilde{\mathbb{P}} \left(Z_t \geq t^{\frac{1}{2}}y \right) \\ &=: J_{31}^{(1)}(t, y) + J_{31}^{(2)}(t, y). \end{aligned} \quad (\text{B.28})$$

By (2.22), it is easy to see that $J_{31}^{(2)}(t, y) \leq \kappa t^{1-\frac{Y}{2}} y^{-Y}$, for any $0 < t \leq 1$ and $y \geq 0$. Hence, dominated convergence theorem together with (2.20) lead to

$$\lim_{t \rightarrow 0} e^{-\tilde{\eta}t} t^{\frac{Y}{2}-1} \int_0^\infty J_{31}^{(2)}(t, y) \frac{1 - e^{\sqrt{t}y}}{\sqrt{t}} \frac{e^{-\frac{y^2}{2\sigma^2}}}{\sqrt{2\pi\sigma^2}} dy = -\frac{C_+}{Y} \int_0^\infty y^{1-Y} \frac{e^{-\frac{y^2}{2\sigma^2}}}{\sqrt{2\pi\sigma^2}} dy. \quad (\text{B.29})$$

As usual, $J_{31}^{(1)}(t, y)$ can be further decomposed as:

$$J_{31}^{(1)}(t, y) = \tilde{\mathbb{E}} \left((e^{-\tilde{U}_t} - 1) \mathbf{1}_{\{Z_t \geq t^{\frac{1}{2}}y, \tilde{U}_t \geq 0\}} \right) + \int_0^\infty e^u \tilde{\mathbb{P}} \left(Z_t \geq t^{\frac{1}{2}}y, \tilde{U}_t \leq -u \right) du.$$

Now, for $u > 0$, $y > 0$ and $t > 0$, by Markov's inequality and (3.2-ii), there exist $\tilde{\kappa} < \infty$ and $0 < t_0 < 1$, such that

$$\tilde{\mathbb{P}} \left(Z_t \geq t^{\frac{1}{2}}y, \tilde{U}_t \leq -u \right) \leq \tilde{\mathbb{P}} \left(\tilde{U}_t \leq -u \right) \leq \tilde{\mathbb{E}} \left(e^{-t^{-1/Y} \tilde{U}_t} \right) e^{-t^{-1/Y} u} \leq \tilde{\kappa} e^{-t^{-1/Y} u}, \quad (\text{B.30})$$

for all $0 < t < t_0$. Also, as we did in (B.24), there exist $K < \infty$ and $0 < t_0 < 1$ such that

$$0 \leq \tilde{\mathbb{E}} \left((1 - e^{-\tilde{U}_t}) \mathbf{1}_{\{Z_t \geq t^{\frac{1}{2}}y, \tilde{U}_t \geq 0\}} \right) \leq \tilde{\mathbb{E}} \left(\tilde{U}_t \right)^+ \leq K t^{1/Y}, \quad (\text{B.31})$$

for all $0 < t < t_0$. Hence, for any $y \geq 0$ and $0 < t < t_0 < 1$, and since $1 - Y/2 < 1/2 < 1/Y$, for $1 < Y < 2$,

$$t^{\frac{Y}{2}-1} |J_{31}^{(1)}(t, y)| \leq \tilde{\kappa} t^{\frac{Y}{2}-1} \int_0^\infty e^{-u(t^{-\frac{1}{Y}}-1)} du + K t^{\frac{1}{Y}+\frac{Y}{2}-1} = \tilde{\kappa} t^{\frac{Y}{2}-1} \frac{t^{\frac{1}{Y}}}{1-t^{\frac{1}{Y}}} + K t^{\frac{1}{Y}+\frac{Y}{2}-1} \rightarrow 0, \quad (\text{B.32})$$

as $t \rightarrow 0$. Since both control functions in (B.30) and (B.31) are independent of y , combining (B.29) and (B.32), and by the dominated convergence theorem,

$$\lim_{t \rightarrow 0} e^{-\tilde{\eta}t} t^{\frac{Y}{2}-2} \int_0^\infty J_{31}(t, y) \frac{1 - e^{\sqrt{t}y}}{\sqrt{t}} \frac{e^{-\frac{y^2}{2\sigma^2}}}{\sqrt{2\pi\sigma^2}} dy = -\frac{C_+}{Y} \int_0^\infty y^{1-Y} \frac{e^{-\frac{y^2}{2\sigma^2}}}{\sqrt{2\pi\sigma^2}} dy. \quad (\text{B.33})$$

Next, by the self-similarity of $(Z_t)_{t \geq 0}$, we decompose the expression $J_{32}(t, y) := \widetilde{\mathbb{E}} \left(e^{-\tilde{U}_t} \mathbf{1}_{\{Z_t \geq t^{\frac{1}{2}} y\}} (1 - e^{-Z_t}) / \sqrt{t} \right)$ appearing in (B.27) as:

$$\begin{aligned} J_{32}(t, y) &= \widetilde{\mathbb{E}} \left(\mathbf{1}_{\{Z_t \geq t^{\frac{1}{2}} y\}} \left(\frac{e^{-\tilde{U}_t} - e^{-(Z_t + \tilde{U}_t)}}{\sqrt{t}} - t^{-\frac{1}{2}} Z_t \right) \right) + \widetilde{\mathbb{E}} \left(t^{\frac{1}{Y} - \frac{1}{2}} Z_1 \mathbf{1}_{\{Z_1 \geq t^{\frac{1}{2} - \frac{1}{Y}} y\}} \right) \\ &=: J_{32}^{(1)}(t, y) + J_{32}^{(2)}(t, y). \end{aligned} \quad (\text{B.34})$$

Note that $J_{32}^{(2)}(t, y)$ is quite similar to $J_{12}(t, y)$ in (B.4) and, thus, the corresponding integral has an asymptotic behavior similar to (B.6). Concretely,

$$\lim_{t \rightarrow 0} t^{\frac{Y}{2} - 1} e^{-\tilde{\eta}t} \int_0^\infty \widetilde{\mathbb{E}} \left(t^{\frac{1}{Y} - \frac{1}{2}} Z_1 \mathbf{1}_{\{Z_1 \geq t^{\frac{1}{2} - \frac{1}{Y}} y\}} \right) e^{\sqrt{t}y} \frac{e^{-\frac{y^2}{2\sigma^2}}}{\sqrt{2\pi\sigma^2}} dy = \frac{C_+}{Y-1} \int_0^\infty y^{1-Y} \frac{e^{-\frac{y^2}{2\sigma^2}}}{\sqrt{2\pi\sigma^2}} dy. \quad (\text{B.35})$$

Next, decompose $J_{32}^{(1)}(t, y)$ as:

$$\begin{aligned} J_{32}^{(1)}(t, y) &= t^{-\frac{1}{2}} \widetilde{\mathbb{E}} \left(\mathbf{1}_{\{Z_t \geq t^{\frac{1}{2}} y\}} \int_{\tilde{U}_t}^{Z_t + \tilde{U}_t} (e^{-x} - 1) dx \right) \\ &= \frac{1}{\sqrt{t}} \int_{-\infty}^0 (e^{-x} - 1) \widetilde{\mathbb{P}} \left(Z_t \geq t^{\frac{1}{2}} y, \tilde{U}_t \leq x \leq Z_t + \tilde{U}_t \right) dx + \frac{1}{\sqrt{t}} \int_0^\infty (e^{-x} - 1) \widetilde{\mathbb{P}} \left(Z_t \geq t^{\frac{1}{2}} y, \tilde{U}_t \leq x \leq Z_t + \tilde{U}_t \right) dx. \end{aligned}$$

Note that for $x > 0$,

$$\widetilde{\mathbb{P}} \left(Z_t \geq t^{\frac{1}{2}} y, \tilde{U}_t \leq x \leq Z_t + \tilde{U}_t \right) \leq \widetilde{\mathbb{P}} \left(x \leq Z_t + \tilde{U}_t \right),$$

while for $x < 0$,

$$\widetilde{\mathbb{P}} \left(Z_t \geq t^{\frac{1}{2}} y, \tilde{U}_t \leq x \leq Z_t + \tilde{U}_t \right) \leq \widetilde{\mathbb{P}} \left(\tilde{U}_t \leq x \right).$$

Using the estimates (B.8) and (B.12), arguments as in getting (B.9) and (B.13) give

$$\lim_{t \rightarrow 0} e^{-\tilde{\eta}t} t^{\frac{Y}{2} - 1} \int_0^\infty J_{32}^{(1)}(t, y) e^{\sqrt{t}y} \frac{e^{-\frac{y^2}{2\sigma^2}}}{\sqrt{2\pi\sigma^2}} dy = 0. \quad (\text{B.36})$$

Combining (B.33), (B.35) and (B.36),

$$\lim_{t \rightarrow 0} t^{\frac{Y}{2} - 1} I_3(t) = \frac{C_+}{Y(Y-1)} \int_0^\infty y^{1-Y} \frac{e^{-\frac{y^2}{2\sigma^2}}}{\sqrt{2\pi\sigma^2}} dy. \quad (\text{B.37})$$

Finally, from (B.2), (B.15), (B.26) and (B.37), and since $1 - Y/2 < 1/2$, for $1 < y < 2$, we obtain (4.2), therefore finishing the proof. $\square$

Proof of Proposition 4.3.

When the diffusion component is nonzero, [43, Proposition 5] implies that $\hat{\sigma}(t) \rightarrow \sigma$ as $t \rightarrow 0$. In particular, $\hat{\sigma}_t t^{1/2} \rightarrow 0$ as $t \rightarrow 0$ and, thus, (A.20) above remains true. Let $\tilde{\sigma}(t) := \hat{\sigma}_t - \sigma$, then $\tilde{\sigma}(t) \rightarrow 0$ as $t \rightarrow 0$, and (A.20) can be written as

$$\begin{aligned} C_{BS}(t, \hat{\sigma}(t)) &= \frac{\sigma}{\sqrt{2\pi}} t^{1/2} + \frac{\tilde{\sigma}(t)}{\sqrt{2\pi}} t^{1/2} - \frac{\hat{\sigma}(t)^3}{24\sqrt{2\pi}} t^{3/2} + O \left((\hat{\sigma}(t) t^{1/2})^5 \right) \\ &= \frac{\sigma}{\sqrt{2\pi}} t^{1/2} + \frac{\tilde{\sigma}(t)}{\sqrt{2\pi}} t^{1/2} + O(t^{3/2}). \end{aligned} \quad (\text{B.38})$$

By comparing (4.3)-(4.4) and (B.38), we have

$$\frac{(C_+ + C_-)2^{-Y}\sigma^{1-Y}}{Y(Y-1)\sqrt{\pi}}\Gamma\left(1 - \frac{Y}{2}\right)t^{\frac{3-Y}{2}} \sim \frac{\tilde{\sigma}(t)}{\sqrt{2\pi}}t^{1/2}, \quad t \rightarrow 0,$$

and, therefore,

$$\tilde{\sigma}(t) \sim \frac{(C_+ + C_-)2^{\frac{1}{2}-Y}\sigma^{1-Y}}{Y(Y-1)}\Gamma\left(1 - \frac{Y}{2}\right)t^{1-\frac{Y}{2}}, \quad t \rightarrow 0.$$

The proof is now complete. $\square$

C Additional proofs

Verification of (2.22).

It suffices to show that, for some constant $\kappa < \infty$ and any $v > 0$ and $0 < t \leq 1$,

$$\tilde{\mathbb{P}}\left(Z_1 \geq t^{-1/Y}v\right) \leq \kappa t|v|^{-Y}.$$

Indeed, using the same argument for $-Z_1$, one can show an analog bound for $\tilde{\mathbb{P}}(Z_1 \leq t^{-1/Y}v)$, with $v < 0$, which in turn will imply (2.22). First, in light of (2.19), there exist $0 < R < \infty$ and $H > 0$, such that for any $u \geq H$,

$$p_Z(u) \leq Ru^{-Y-1}. \quad (\text{C.1})$$

Thus, whenever $t^{-1/Y}v \geq H$,

$$\tilde{\mathbb{P}}\left(Z_1 \geq t^{-1/Y}v\right) \leq R \int_{t^{-1/Y}v}^{\infty} u^{-Y-1} du = \frac{R}{Y} \left(t^{-1/Y}v\right)^{-Y} = \frac{R}{Y} tv^{-Y}.$$

Since $\tilde{\mathbb{P}}(Z_1 \geq t^{-1/Y}v) \leq 1 < H^Y tv^{-Y}$, when $t^{-1/Y}v < H$, we deduce that

$$\begin{aligned} \tilde{\mathbb{P}}\left(Z_1 \geq t^{-1/Y}v\right) &\leq \mathbf{1}_{\{t^{-1/Y}v \geq H\}} \tilde{\mathbb{P}}\left(Z_1 \geq t^{-1/Y}v\right) + \mathbf{1}_{\{t^{-1/Y}v < H\}} \tilde{\mathbb{P}}\left(Z_1 \geq t^{-1/Y}v\right) \\ &\leq \mathbf{1}_{\{t^{-1/Y}v \geq H\}} \frac{R}{Y} tv^{-Y} + \mathbf{1}_{\{t^{-1/Y}v < H\}} H^Y tv^{-Y} \\ &\leq tv^{-Y} \left(\frac{R}{Y} + H^Y \right), \end{aligned}$$

for all $0 < t \leq 1$ and $v > 0$. $\square$

Verification of (B.5)

This is similar to the proof of (2.22). Using (C.1), for any $t > 0$ and $y > 0$,

$$\begin{aligned} t^{\frac{Y}{2}-1} J_{12}(t, y) &= t^{\frac{Y}{2}-1} t^{\frac{1}{Y}-\frac{1}{2}} \int_{t^{\frac{1}{2}-\frac{1}{Y}}y}^{\infty} up_Z(u) du \\ &= t^{(\frac{1}{Y}-\frac{1}{2})(1-Y)} \mathbf{1}_{\{t^{\frac{1}{2}-\frac{1}{Y}}y \geq H\}} \int_{t^{\frac{1}{2}-\frac{1}{Y}}y}^{\infty} up_Z(u) du \\ &\quad + t^{(\frac{1}{Y}-\frac{1}{2})(1-Y)} \mathbf{1}_{\{t^{\frac{1}{2}-\frac{1}{Y}}y \leq H\}} \left(\int_H^{\infty} up_Z(u) du + \int_{t^{\frac{1}{2}-\frac{1}{Y}}y}^H up_Z(u) du \right) \\ &\leq t^{(\frac{1}{Y}-\frac{1}{2})(1-Y)} \mathbf{1}_{\{t^{\frac{1}{2}-\frac{1}{Y}}y \geq H\}} \int_{t^{\frac{1}{2}-\frac{1}{Y}}y}^{\infty} Ru^{-Y} du \\ &\quad + t^{(\frac{1}{Y}-\frac{1}{2})(1-Y)} \mathbf{1}_{\{t^{\frac{1}{2}-\frac{1}{Y}}y \leq H\}} \left(\int_H^{\infty} Ru^{-Y} du + H \tilde{\mathbb{P}}(Z_1 \geq t^{\frac{1}{2}-\frac{1}{Y}}y) \right). \end{aligned}$$

Then, using (2.22),

$$\begin{aligned}
t^{\frac{Y}{2}-1} J_{12}(t, y) &\leq t^{(\frac{1}{Y}-\frac{1}{2})(1-Y)} R(Y-1)^{-1} y^{1-Y} t^{(\frac{1}{2}-\frac{1}{Y})(1-Y)} \\
&\quad + t^{(\frac{1}{Y}-\frac{1}{2})(1-Y)} \mathbf{1}_{\{t^{\frac{1}{2}-\frac{1}{Y}} y \leq H\}} \left(R H^{1-Y} + H \left(\frac{H}{t^{\frac{1}{2}-\frac{1}{Y}} y} \right)^{Y-1} \right) \\
&\leq R(Y-1)^{-1} y^{1-Y} + t^{(\frac{1}{Y}-\frac{1}{2})(1-Y)} R t^{(\frac{1}{2}-\frac{1}{Y})(1-Y)} y^{1-Y} + H^Y y^{1-Y},
\end{aligned}$$

and (B.5) follows with $\lambda := RY/(Y-1) + H^Y$. □

Acknowledgments: The authors gratefully acknowledge the insightful comments provided by an Associate Editor and two anonymous referees, which significantly contributed to improve the quality of the manuscript.

References

- [1] O. E. Barndorff-Nielsen, Processes of Normal Inverse Gaussian type, *Finance and Stochastics*, 2, 41-68, 1997.
- [2] H. Berestycki, J. Busca, and I. Florent. Asymptotics and calibration of local volatility models, *Quantitative Finance*, 2, 61-69, 2002.
- [3] H. Berestycki, J. Busca, and I. Florent. Computing the implied volatility in stochastic volatility models, *Communications on Pure and Applied Mathematics*, Vol LVII, 1352-1373, 2004.
- [4] Boyarchenko, S.I., and S.Z. Levendorskii. Non-Gaussian Merton-Black- Scholes theory, *Adv. Ser. Stat. Sci. Appl. Probab.* 9. World Scientific Publishing Co., Inc., River Edge, NJ, 2002.
- [5] P. Carr, H. Geman, D. Madan, and M. Yor. The fine structure of asset returns: an empirical investigation, *Journal of Business*, 75, 305-332, 2002.
- [6] P. Carr and D. Madan. Saddle point methods for option pricing, *The Journal of Computational Finance*, 13(1), 49-61, 2009.
- [7] R. Cont, J. Bouchaud, and M. Potters. Scaling in financial data: stable laws and beyond, in *Scale Invariance and Beyond*, Dubrulle, B., Graner, F., and Sornette, D., eds., 1997.
- [8] R. Cont and P. Tankov, *Financial modelling with jump processes*, Chapman & Hall, 2004.
- [9] P. Carr and L. Wu. What type of process underlies options? A simple robust test, *Journal of Finance*, 58(6), 2581-2610, 2003.
- [10] E. Eberlein, U. Keller, and K. Prause. New insights into smile, mispricing and value at risk, *Journal of Bussiness*, 71, 371-406, 1998.
- [11] J. Feng, M. Forde, and J.P. Fouque. Short maturity asymptotics for a fast mean reverting Heston stochastic volatility model, *SIAM Journal on Financial Mathematics*, 1, 126-141, 2010.
- [12] J. Feng, J.P. Fouque, and R. Kumar, Small-time Asymptotics for Fast Mean-Reverting Stochastic Volatility Models, Forthcoming in *The Annals of Applied Probability*, 2012.
- [13] J.E. Figueroa-López and M. Forde. The small-maturity smile for exponential Lévy models, *SIAM Journal on Financial Mathematics* 3(1), 33-65, 2012.
- [14] J.E. Figueroa-López and C. Houdré. Small-time expansions for the transition distributions of Lévy processes, *Stochastic Processes and their Applications*, 119, 3862-3889, 2009.

- [15] J.E. Figueroa-López, R. Gong, and C. Houdré. Small-time expansions of the distributions, densities, and option prices under stochastic volatility models with Lévy jumps, *Stochastic Processes and their Applications*, 122, 1808–1839, 2012.
- [16] J.E. Figueroa-López, R. Gong, and C. Houdré. High-order short-time expansions for ATM option prices under the CGMY model, *Preprint. Available at arXiv:1112.3111v1 [q-fin.CP]*.
- [17] M. Forde and A. Jacquier. Small-time asymptotics for implied volatility under the Heston model, *Int. J. Theor. Appl. Finance*, 12(6), 861–876, 2009.
- [18] M. Forde and A. Jacquier. Small time asymptotics for an uncorrelated local-stochastic volatility model, *Applied Mathematical Finance*, 18(6), 517–535, 2011.
- [19] M. Forde, A. Jacquier, and R. Lee. The small-time smile and term structure for implied volatility under the Heston model, *preprint*, 2011.
- [20] J. Gatheral, E. Hsu, P. Laurence, C. Ouyang, and T-H. Wang. Asymptotics of implied volatility in local volatility models, Forthcoming in *Mathematical Finance*, 2012.
- [21] P. Henry-Labordère. Analysis, geometry, and modeling in finance: advanced methods in option pricing, *Chapman & Hall*, 2009.
- [22] C. Houdré. Remarks on deviation inequalities for functions of infinitely divisible random vectors, *Annals of Probability*, 30(3), 1223–1237, 2002.
- [23] O. Kallenberg. *Foundations of Modern Probability*. Springer-Verlag, Berlin, New York, Heidelberg, 1997.
- [24] I. Koponen. Analytic approach to the problem of convergence of truncated Lévy flights towards the Gaussian stochastic process, *Physical Review E*, 52, 1197–1199, 1995.
- [25] S. Kou. A jump-diffusion model for option pricing, *Management Science*, 48, 1086–1101, 2002.
- [26] D. B. Madan, P. Carr, and E. Chang. The variance gamma process and option pricing, *European Finance Review*, 2, 79–105, 1998.
- [27] D. B. Madan and F. Milne. Option pricing with VG martingale components, *Mathematical Finance*, 1, 39–56, 1991.
- [28] D. B. Madan and E. Seneta. The variance gamma (VG) model for share market returns, *Journal of Business*, 63, 511–524, 1990.
- [29] B. Mandelbrot. The variation of certain speculative prices. *The Journal of Business*, 36:394–419, 1963.
- [30] A. Matacz. Financial modeling and option theory with the truncated Lévy process, *Int. J. Theor. Appl. Finance*, 3:143–160, 2000.
- [31] R. Merton. Option pricing when underlying stock returns are discontinuous, *Journal of Financial Economics*, 3, 125–144, 1976.
- [32] J. Muhle-Karbe and M. Nutz. Small-time asymptotics of option prices and first absolute moments, *Journal of Applied Probability*, 48(4), 1003–1020, 2011.
- [33] L. Paulot. Asymptotic implied volatility at the second order with application to the SABR model, *Preprint*, 2009.
- [34] S.J. Press. A compound event model for security prices. *The Journal of Business*, 40:317–335, 1967.
- [35] M. Roper. Implied volatility: small time to expiry asymptotics in exponential Lévy models, *Thesis, University of New South Wales*, 2009.
- [36] M. Roper and M. Rutkowski. A note on the behaviour of the Black-Scholes implied volatility close to expiry, *preprint*, 2007.

- [37] M. Rosenbaum and P. Tankov. Asymptotic results for time-changed Lévy processes sampled at hitting times. *Stochastic processes and their applications*, 121:1607–1633, 2011.
- [38] L. Rüschendorf and J. Woerner. Expansion of transition distributions of Lévy processes in small time. *Bernoulli*, 8, 81-96, 2002.
- [39] J. Rosiński. Tempering stable processes. *Stochastic processes and their applications*, 117:677–707, 2007.
- [40] J. Rosiński and J. L. Sinclair. Generalized tempered stable processes, *Banach Center Publication*, 90, 153-170, 2010.
- [41] G. Samorodnitsky and M. Taqqu. *Stable non-Gaussian random processes*, Chapman & Hall, New York, 1994.
- [42] K. Sato. *Lévy processes and infinitely divisible distributions*, Cambridge University Press, 1999.
- [43] P. Tankov. Pricing and hedging in exponential Lévy models: review of recent results, *Paris-Princeton Lecture Notes in Mathematical Finance*, Springer 2010.